

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
6 September 2002 (06.09.2002)

PCT

(10) International Publication Number  
WO 02/068442 A1

(51) International Patent Classification<sup>7</sup>: C07H 19/01, A01N 43/90

(21) International Application Number: PCT/EP02/02044

(22) International Filing Date: 26 February 2002 (26.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0373/01 27 February 2001 (27.02.2001) CH

(71) Applicant (*for all designated States except US*): SYNGENTA PARTICIPATIONS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): PITTERNA, Thomas [AT/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH).

(74) Agent: BASTIAN, Werner; Syngenta Participations AG, Intellectual Property, P.O. Box, CH-4002 Basel (CH).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: SALTS OF AVERMECTINS SUBSTITUTED IN THE 4"-POSITION AND HAVING PESTICIDAL PROPERTIES


(57) Abstract: A compound of formula wherein X<sup>-</sup> is an anion; n is 1, 2, 3 or 4; R<sub>1</sub> is C<sub>1</sub>-C<sub>12</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl; or C<sub>2</sub>-C<sub>12</sub>alkenyl; R<sub>2</sub> is hydrogen, unsubstituted or substituted C<sub>1</sub>-C<sub>12</sub>alkyl or C<sub>2</sub>-C<sub>12</sub>alkenyl; R<sub>3</sub> is hydrogen, unsubstituted or substituted C<sub>1</sub>-C<sub>12</sub>alkyl, C<sub>3</sub>-C<sub>12</sub>cycloalkyl, C<sub>2</sub>-C<sub>12</sub>alkenyl; or C<sub>2</sub>-C<sub>12</sub>alkynyl; or R<sub>2</sub> and R<sub>3</sub> together are an alkylene or alkenylene bridge; and, where applicable, E/Z isomers, E/Z isomeric mixtures and/or tautomers, with the proviso that R<sub>1</sub> is not sec-butyl or isopropyl when R<sub>2</sub> is H and R<sub>3</sub> is methyl; and, where applicable, their possible tautomers; a process for the preparation of those compounds and their tautomers and the use thereof; pesticidal compositions in which the active ingredient has been selected from those compounds and their tautomers; and a process for the preparation of those compositions and the use thereof; intermediates, in free form or in salt form, for the preparation of those compounds and, where applicable, their tautomers, in free form or in salt form, are described.



WO 02/068442 A1

Salts of avermectins substituted in the 4"-position and having pesticidal properties

The invention relates to (1) a compound of formula



wherein

$X^-$  is an anion;

$n$  is 1, 2, 3 or 4;

$R_1$  is  $C_1$ - $C_{12}$ alkyl,  $C_3$ - $C_8$ cycloalkyl; or  $C_2$ - $C_{12}$ alkenyl;

$R_2$  is hydrogen, unsubstituted or mono- to penta-substituted  $C_1$ - $C_{12}$ alkyl, unsubstituted or mono- to penta-substituted  $C_2$ - $C_{12}$ alkenyl;

$R_3$  is hydrogen, unsubstituted or mono- to penta-substituted  $C_1$ - $C_{12}$ alkyl, unsubstituted or mono- to penta-substituted  $C_3$ - $C_{12}$ cycloalkyl, unsubstituted or mono- to penta-substituted  $C_2$ - $C_{12}$ alkenyl; unsubstituted or mono- to penta-substituted  $C_2$ - $C_{12}$ alkynyl; or

$R_2$  and  $R_3$  together are a three- to seven-membered alkylene bridge, or a four- to seven-membered alkenylene bridge wherein a  $-CH_2-$  group may have been replaced by O, S or  $NR_4$ ;

and wherein the substituents of the mentioned alkyl, alkenyl, alkynyl, alkylene, alkenylene and cycloalkyl radicals are selected from the group consisting of OH, halogen, halo- $C_1$ - $C_2$ alkyl, CN,  $NO_2$ ,  $C_2$ - $C_6$ alkynyl,  $C_3$ - $C_8$ cycloalkyl, norbornylenyl,  $C_3$ - $C_8$ cycloalkenyl;  $C_3$ - $C_8$ cycloalkenyl unsubstituted or substituted by from one to three methyl groups;

C<sub>3</sub>-C<sub>8</sub>halocycloalkyl, C<sub>1</sub>-C<sub>12</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>cycloalkoxy, C<sub>1</sub>-C<sub>12</sub>haloalkoxy, C<sub>1</sub>-C<sub>12</sub>alkylthio, C<sub>3</sub>-C<sub>8</sub>cycloalkylthio, C<sub>1</sub>-C<sub>12</sub>haloalkylthio, C<sub>1</sub>-C<sub>12</sub>alkylsulfinyl, C<sub>3</sub>-C<sub>8</sub>cycloalkylsulfinyl, C<sub>1</sub>-C<sub>12</sub>haloalkylsulfinyl, C<sub>3</sub>-C<sub>8</sub>halocycloalkylsulfinyl, C<sub>1</sub>-C<sub>12</sub>alkylsulfonyl, C<sub>3</sub>-C<sub>8</sub>cycloalkylsulfonyl, C<sub>1</sub>-C<sub>12</sub>haloalkylsulfonyl, C<sub>3</sub>-C<sub>8</sub>halocycloalkylsulfonyl, C<sub>2</sub>-C<sub>8</sub>alkenyl, C<sub>2</sub>-C<sub>8</sub>alkynyl, NH(C<sub>1</sub>-C<sub>6</sub>alkyl), N(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub>, -C(=O)R<sub>5</sub>, -NHC(=O)R<sub>6</sub>, -P(=O)(OC<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub>;

aryl, heterocyclyl, aryloxy, heterocyclyloxy; and also aryl, heterocyclyl, aryloxy and heterocyclyloxy that, depending upon the possibilities of substitution at the ring, are mono- to penta-substituted by substituents selected from the group consisting of OH, halogen, CN, NO<sub>2</sub>, C<sub>1</sub>-C<sub>12</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>1</sub>-C<sub>12</sub>haloalkyl, C<sub>1</sub>-C<sub>12</sub>alkoxy, C<sub>1</sub>-C<sub>12</sub>haloalkoxy, C<sub>1</sub>-C<sub>12</sub>alkylthio, C<sub>1</sub>-C<sub>12</sub>haloalkylthio, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, dimethylamino-C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>2</sub>-C<sub>8</sub>alkenyl, C<sub>2</sub>-C<sub>8</sub>alkynyl, phenoxy, phenyl-C<sub>1</sub>-C<sub>6</sub>alkyl; phenoxy unsubstituted or substituted by from one to three substituents selected independently of one another from halogen, methoxy, trifluoromethyl and trifluoromethoxy; phenyl-C<sub>1</sub>-C<sub>6</sub>alkoxy unsubstituted or substituted in the aromatic ring by from one to three substituents selected independently of one another from halogen, methoxy, trifluoromethyl and trifluoromethoxy; phenyl-C<sub>2</sub>-C<sub>6</sub>alkenyl, phenyl-C<sub>2</sub>-C<sub>6</sub>alkynyl, methylenedioxy, -C(=O)R<sub>5</sub>, -O-C(=O)R<sub>6</sub>, -NH-C(=O)R<sub>6</sub>, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>12</sub>alkyl), N(C<sub>1</sub>-C<sub>12</sub>alkyl)<sub>2</sub>, C<sub>1</sub>-C<sub>6</sub>alkylsulfinyl, C<sub>3</sub>-C<sub>8</sub>cycloalkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>haloalkylsulfinyl, C<sub>3</sub>-C<sub>8</sub>halocycloalkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>alkylsulfonyl, C<sub>3</sub>-C<sub>8</sub>cycloalkylsulfonyl, C<sub>1</sub>-C<sub>6</sub>haloalkylsulfonyl and C<sub>3</sub>-C<sub>8</sub>halocycloalkylsulfonyl;

R<sub>4</sub> is C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>2</sub>-C<sub>8</sub>alkenyl, C<sub>2</sub>-C<sub>8</sub>alkynyl, benzyl or -C(=O)-R<sub>5</sub>;

R<sub>5</sub> is H, OH, SH, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>12</sub>alkyl), N(C<sub>1</sub>-C<sub>12</sub>alkyl)<sub>2</sub>, C<sub>1</sub>-C<sub>12</sub>alkyl, C<sub>1</sub>-C<sub>12</sub>haloalkyl, C<sub>1</sub>-C<sub>12</sub>alkoxy, C<sub>1</sub>-C<sub>12</sub>haloalkoxy, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>12</sub>alkylthio, C<sub>2</sub>-C<sub>8</sub>alkenyloxy, C<sub>2</sub>-C<sub>8</sub>alkynyloxy; phenyl, phenoxy, benzyloxy, -NH-phenyl, -N(C<sub>1</sub>-C<sub>6</sub>alkyl)-phenyl, NH-C<sub>1</sub>-C<sub>6</sub>alkyl-C(=O)-R<sub>7</sub>, -N(C<sub>1</sub>-C<sub>6</sub>alkyl)-C<sub>1</sub>-C<sub>6</sub>alkyl-C(=O)-R<sub>7</sub>; or phenyl, phenoxy, benzyloxy, NH-phenyl or -N(C<sub>1</sub>-C<sub>6</sub>alkyl)-phenyl substituted in the aromatic ring by from one to three substituents selected independently of one another from halogen, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>haloalkyl and C<sub>1</sub>-C<sub>6</sub>haloalkoxy;

R<sub>6</sub> is H, C<sub>1</sub>-C<sub>12</sub>alkyl, C<sub>1</sub>-C<sub>12</sub>haloalkyl, C<sub>2</sub>-C<sub>8</sub>alkenyl, C<sub>2</sub>-C<sub>8</sub>alkynyl, phenyl, benzyl, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>12</sub>alkyl), N(C<sub>1</sub>-C<sub>12</sub>alkyl)<sub>2</sub>, -NH-phenyl or -N(C<sub>1</sub>-C<sub>12</sub>alkyl)-phenyl; and

R<sub>7</sub> is H, OH, C<sub>1</sub>-C<sub>12</sub>alkyl, C<sub>1</sub>-C<sub>12</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>2</sub>-C<sub>8</sub>alkenyloxy, phenyl, phenoxy, benzyloxy, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>12</sub>alkyl), N(C<sub>1</sub>-C<sub>12</sub>alkyl)<sub>2</sub>, -NH-phenyl or -N(C<sub>1</sub>-C<sub>12</sub>alkyl)-phenyl;

and, where applicable, to E/Z isomers, E/Z isomeric mixtures and/or tautomers;

with the proviso that R<sub>1</sub> is not sec-butyl or isopropyl when R<sub>2</sub> is H and R<sub>3</sub> is methyl;

to a process for the preparation of those compounds and their isomers and tautomers and to the use thereof; to pesticidal compositions in which the active ingredient has been selected from those compounds and their tautomers; and to a method of controlling pests using those compositions.

Certain macrolide compounds are proposed for pest control in the literature, for example in USP-4 427 663. The biological properties of those known compounds are not entirely satisfactory, however, for which reason there is a need to provide further compounds having pesticidal properties, especially for the control of insects and members of the order Acarina. That problem is solved according to the invention by the provision of the present compounds of formula (I).

The compounds claimed according to the invention are derivatives of avermectin. Avermectins are known to the person skilled in the art. They are a group of structurally closely related pesticidally active compounds which are obtained by fermentation of a strain of the microorganism Streptomyces avermitilis. Derivatives of avermectins can be obtained via conventional chemical syntheses.

The avermectins obtainable from Streptomyces avermitilis are designated A1a, A1b, A2a, A2b, B1a, B1b, B2a and B2b. Compounds with the designation "A" have a methoxy radical in the 5-position; those compounds designated "B" have an OH group. The "a" series comprises compounds wherein the substituent R<sub>1</sub> (in position 25) is a sec-butyl radical; the "b" series have an isopropyl radical in the 25-position. The number 1 in the name of a compound indicates that atoms 22 and 23 are bonded by a double bond; the number 2 indicates that they are bonded by a single bond and carbon atom 23 carries an OH group. The above designations are retained in the present Application in order in the case of the non-natural avermectin derivatives according to the invention to indicate the specific structural type, which corresponds to natural avermectin. There are claimed according to the invention salts of compounds of the B1 series, more especially mixtures of salts of avermectin derivatives B1a and B1b.

Some of the compounds of formula (I) may be in the form of tautomers. Accordingly, any reference to the compounds of formula (I) hereinabove and hereinbelow is to be under-

stood, where applicable, as including also corresponding tautomers, even if the latter are not specifically mentioned in every case.

The general terms used hereinabove and hereinbelow have the meanings given below, unless defined to the contrary.

Unless defined otherwise, carbon-containing groups and compounds each contain from 1 up to and including 6, preferably from 1 up to and including 4, especially 1 or 2, carbon atoms.

Halogen - as a group *per se* and as a structural element of other groups and compounds, such as haloalkyl, haloalkoxy and haloalkylthio - is fluorine, chlorine, bromine or iodine, especially fluorine, chlorine or bromine, more especially fluorine or chlorine. In the cases where halogen functions as a leaving group, bromine and iodine are preferred.

Alkyl - as a group *per se* and as a structural element of other groups and compounds, such as haloalkyl, alkoxy and alkylthio - is, in each case giving due consideration to the number of carbon atoms contained in the group or compound in question, either straight-chained, i.e. methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl or octyl, or branched, for example isopropyl, isobutyl, sec-butyl, tert-butyl, isopentyl, neopentyl or isohexyl.

Cycloalkyl - as a group *per se* and as a structural element of other groups and compounds, such as halocycloalkyl, cycloalkoxy and cycloalkylthio - is, in each case giving due consideration to the number of carbon atoms contained in the group or compound in question, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl.

Alkenyl - as a group *per se* and as a structural element of other groups and compounds - is, in each case giving due consideration to the number of carbon atoms and conjugated or isolated double bonds contained in the group in question, either straight-chained, e.g. allyl, 2-butenyl, 3-pentenyl, 1-hexenyl, 1-heptenyl, 1,3-hexadienyl or 1,3-octadienyl, or branched, e.g. isopropenyl, isobutenyl, isoprenyl, tert-pentenyl, isohexenyl, isoheptenyl or isooctenyl. Alkenyl groups having from 3 to 12, especially from 3 to 6, more especially 3 or 4, carbon atoms are preferred.

Alkynyl - as a group *per se* and as a structural element of other groups and compounds - is, in each case giving due consideration to the number of carbon atoms and conjugated or isolated double bonds contained in the group or compound in question, either straight-chained, e.g. propargyl, 2-butyne, 3-pentyne, 1-hexyne, 1-heptyne, 3-hexen-1-yne or 1,5-heptadien-3-yne, or branched, e.g. 3-methylbut-1-yne, 4-ethylpent-1-yne, 4-methyl-

hex-2-ynyl or 2-methylhept-3-ynyl. Alkynyl groups having from 3 to 12, especially from 3 to 6, more especially 3 or 4, carbon atoms are preferred.

Alkylene and alkenylene are straight-chained or branched bridge members, especially  $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ ,  $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ ,  $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ ,  $-\text{CH}_2(\text{CH}_3)\text{CH}_2-\text{CH}_2-$ ,  $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{CH}_2-$ ,  $-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-$  or  $-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-\text{CH}_2-$ .

Halo-substituted carbon-containing groups and compounds, such as haloalkyl, haloalkoxy and haloalkylthio, may be partially halogenated or perhalogenated, the halogen substituents in the case of polyhalogenation being the same or different. Examples of haloalkyl - as a group *per se* and as a structural element of other groups and compounds, such as haloalkoxy and haloalkylthio - are methyl substituted from one to three times by fluorine, chlorine and/or bromine, such as  $\text{CHF}_2$  or  $\text{CF}_3$ ; ethyl substituted from one to five times by fluorine, chlorine and/or bromine, such as  $\text{CH}_2\text{CF}_3$ ,  $\text{CF}_2\text{CF}_3$ ,  $\text{CF}_2\text{CCl}_3$ ,  $\text{CF}_2\text{CHCl}_2$ ,  $\text{CF}_2\text{CHF}_2$ ,  $\text{CF}_2\text{CFCl}_2$ ,  $\text{CF}_2\text{CHBr}_2$ ,  $\text{CF}_2\text{CHClF}$ ,  $\text{CF}_2\text{CHBrF}$  or  $\text{CClFCHClF}$ ; propyl or isopropyl substituted from one to seven times by fluorine, chlorine and/or bromine, such as  $\text{CH}_2\text{CHBrCH}_2\text{Br}$ ,  $\text{CF}_2\text{CHF}_2\text{CF}_3$ ,  $\text{CH}_2\text{CF}_2\text{CF}_3$  or  $\text{CH}(\text{CF}_3)_2$ ; butyl or an isomer thereof substituted from one to nine times by fluorine, chlorine and/or bromine, such as  $\text{CF}(\text{CF}_3)\text{CH}_2\text{CF}_3$  or  $\text{CH}_2(\text{CF}_2)_2\text{CF}_3$ ; pentyl or an isomer thereof substituted from one to eleven times by fluorine, chlorine and/or bromine, such as  $\text{CF}(\text{CF}_3)(\text{CHF})_2\text{CF}_3$  or  $\text{CH}_2(\text{CF}_2)_3\text{CF}_3$ ; and hexyl or an isomer thereof substituted from one to thirteen times by fluorine, chlorine and/or bromine, such as  $(\text{CH}_2)_4\text{CHBrCH}_2\text{Br}$ ,  $\text{CF}_2(\text{CHF})_4\text{CF}_3$ ,  $\text{CH}_2(\text{CF}_2)_4\text{CF}_3$  or  $\text{C}(\text{CF}_3)_2(\text{CHF})_2\text{CF}_3$ .

Aryl is especially phenyl, naphthyl, anthracenyl or perylenyl, preferably phenyl.

Heterocyclyl is especially pyridyl, pyrimidyl, s-triazinyl, 1,2,4-triazinyl, thienyl, furyl, tetrahydrofuranyl, pyranal, tetrahydropyranyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, triazolyl, oxazolyl, thiadiazolyl, oxadiazolyl, benzothienyl, quinolynyl, quinoxalynyl, benzofuranyl, benzimidazolyl, benzopyrrolyl, benzothiazolyl, indolyl, coumarinyl or indazolyl, which are preferably bonded *via* a carbon atom; preference is given to thienyl, thiazolyl, benzofuranyl, benzothiazolyl, furyl, tetrahydropyranyl and indolyl; especially pyridyl or thiazolyl.

$\text{X}^-$  is the anion of an inorganic acid, especially a mineral acid, e.g. sulfuric acid, a phosphoric acid or a hydrohalic acid;

the anion of an organic carboxylic acid, such as an unsubstituted or substituted, e.g. halo-substituted,  $\text{C}_1$ - $\text{C}_4$ alkanecarboxylic acid, for example acetic acid, a saturated or unsaturated dicarboxylic acid, for example oxalic acid, malonic acid, maleic acid, fumaric

acid or phthalic acid, a hydroxycarboxylic acid, for example ascorbic acid, lactic acid, malic acid, tartaric acid or citric acid, or benzoic acid;

the anion of an organic sulfonic acid, such as an unsubstituted or substituted, e.g. halo-substituted, C<sub>1</sub>-C<sub>4</sub>alkane- or aryl-sulfonic acid, for example methane- or p-toluene-sulfonic acid;

the anion of an active-H-C compound. Active-H-C compounds include especially organic compounds that carry strongly electron-attracting substituents, such as nitriles, carbonyls or nitro groups. Special preference is given to anions of compounds of formula Y<sub>1</sub>-CH<sub>2</sub>-Y<sub>2</sub> wherein Y<sub>1</sub> and Y<sub>2</sub> denote an electron-attracting group. Special preference is given to the anions of malodinitrile, cyanoacetic acid, esters of cyanoacetic acid, amides of cyanoacetic acid, acetoacetic acid, esters of acetoacetic acid, acetylacetone, cyanacetone and barbituric acid; or

the anion of an acidic phenol, for example picric acid.

Within the scope of the present invention, special preference is given to

(2) compounds according to (1) of formula (I) wherein R<sub>1</sub> is isopropyl or sec-butyl, preferably wherein a mixture of the isopropyl and the sec-butyl derivative is present;

(3) compounds according to one of groups (1) and (2) of formula (I) wherein R<sub>2</sub> is H;

(4) compounds according to one of groups (1) and (2) of formula (I) wherein R<sub>2</sub> is C<sub>1</sub>-C<sub>8</sub>alkyl, especially methyl;

(5) compounds according to one of groups (1), (2) and (4) of formula (I) wherein R<sub>2</sub> is ethyl;

(6) compounds according to one of groups (1), (2) and (4) of formula (I) wherein R<sub>2</sub> is n-propyl;

(7) compounds according to one of groups (1) to (6) of formula (I) wherein R<sub>3</sub> is H or unsubstituted or substituted, especially unsubstituted, C<sub>1</sub>-C<sub>5</sub>alkyl;

(8) compounds according to one of groups (1) to (7) of formula (I) wherein R<sub>3</sub> is methyl;

(9) compounds according to one of groups (1) to (7) of formula (I) wherein R<sub>3</sub> is ethyl;

(10) compounds according to one of groups (1) to (7) of formula (I) wherein R<sub>3</sub> is n-propyl;

(11) compounds according to one of groups (2) to (7) of formula (I) wherein  $R_3$  is isopropyl;

(12) compounds according to one of groups (1) to (7) of formula (I) wherein  $R_3$  is n-butyl, sec-butyl, isobutyl or tert-butyl;

(13) compounds according to one of groups (1) to (6) of formula (I) wherein  $R_3$  is unsubstituted or substituted, especially unsubstituted,  $C_6$ - $C_{12}$ alkyl;

(14) compounds according to one of groups (1) and (2) of formula (I) wherein  $R_2$  and  $R_3$  together are  $-CH_2-CH_2-CH_2-$  or  $-CH_2-CH_2-CH_2-CH_2-$ ;

(15) compounds according to one of groups (1) and (2) of formula (I) wherein  $R_2$  and  $R_3$  together are  $-CH_2-CH_2-O-CH_2-CH_2-$  or  $-CH_2-CH_2-N(CH_3)-CH_2-CH_2-$ ;

(16) compounds according to one of groups (1) to (6) of formula (I) wherein  $R_3$  is substituted  $C_1$ - $C_4$ alkyl and the substituents are selected from the group consisting of OH, halogen,  $C_2$ - $C_8$ alkynyl,  $C_3$ - $C_8$ cycloalkyl;  $C_3$ - $C_8$ cycloalkenyl unsubstituted or substituted by from one to three methyl groups;  $C_1$ - $C_{12}$ alkoxy,  $-C(=O)R_5$ ,  $-NHC(=O)R_6$ ,  $-P(=O)-(OC_1-C_6alkyl)_2$ ; and phenyl, naphthyl, anthracenyl, phenanthrenyl, fluorenyl, perylenyl and heterocyclyl which are unsubstituted or, depending upon the possibilities of substitution at the ring, mono- to penta-substituted;

especially wherein the substituents of  $R_3$  are selected from the group consisting of halogen,  $C_3$ - $C_8$ cycloalkyl,  $C_2$ - $C_8$ alkynyl,  $-C(=O)R_5$ ,  $-NHC(=O)R_6$ ,  $-P(=O)(OC_1-C_6alkyl)_2$ ; and phenyl, naphthyl, anthracenyl, pyridyl, thiazolyl, imidazolyl, furyl, quinolinyl and pyrazolyl which are unsubstituted or, depending upon the possibilities of substitution at the ring, mono- to tri-substituted;

(17) compounds according to one of groups (1) to (6) of formula (I) wherein  $R_3$  is benzyl that carries on the aromatic moiety one to three substituents that are selected from the group consisting of OH, halogen, CN,  $NO_2$ ,  $C_1$ - $C_2$ alkyl, dimethylamino- $C_1$ - $C_4$ alkoxy,  $C_3$ - $C_6$ cycloalkyl,  $C_1$ - $C_2$ haloalkyl,  $C_1$ - $C_2$ alkoxy,  $C_1$ - $C_2$ haloalkoxy, phenoxy, phenyl- $C_1$ - $C_6$ alkyl, phenyl- $C_1$ - $C_4$ alkenyl; phenoxy unsubstituted or substituted by chlorine or methoxy; benzyloxy unsubstituted or substituted by chlorine, methoxy or trifluoromethyl; methylenedioxy,  $-C(=O)R_5$ ,  $-O-C(=O)R_6$  and  $NHC(=O)R_6$ ;

$R_5$  is H, OH,  $NH_2$ ,  $NH(C_1-C_2alkyl)$ ,  $N(C_1-C_2alkyl)_2$ ,  $-O-C_1-C_2alkyl-C(=O)-R_7$ ,  $NHC_1-C_2alkyl-C(=O)-R_7$ ,  $C_1-C_6alkyl$ ,  $C_1-C_2alkoxy$ ,  $C_1-C_2alkoxy-C_1-C_2alkoxy$ ,  $C_2-C_4alkenyloxy$ ,  $C_2-C_4alkynyloxy$ ; phenyl, phenoxy, benzyloxy, NH-phenyl,  $NH-C_1-C_6alkyl-C(=O)-R_7$ ; or



phenyl, phenoxy, benzyloxy or NH-phenyl substituted by halogen, nitro, methoxy, trifluoromethyl or trifluoromethoxy;

$R_6$  is H,  $C_1$ - $C_3$ alkyl, phenyl or benzyl; and

$R_7$  is H, OH,  $NH_2$ ,  $NH(C_1-C_{12}alkyl)$ ,  $N(C_1-C_{12}alkyl)_2$ ,  $C_1$ - $C_{12}alkyl$ ,  $C_1$ - $C_{12}alkoxy$ ,  $C_1$ - $C_6alkoxy$ - $C_1$ - $C_6alkoxy$ ,  $C_2$ - $C_8alkenyloxy$ , phenyl, phenoxy, benzyloxy or NH-phenyl;

(18) compounds according to one of groups (1) to (6) of formula (I) wherein  $R_3$  is  $C_1$ - $C_4alkyl$ - $C(=O)R_5$ , especially  $-CH_2-C(=O)R_5$ ; and

$R_5$  is H, OH,  $NH_2$ ,  $NH(C_1-C_2alkyl)$ ,  $N(C_1-C_2alkyl)_2$ ,  $C_1$ - $C_4alkyl$ ,  $C_1$ - $C_{12}alkoxy$ ,  $C_2$ - $C_4alkenyloxy$ , phenyl, phenoxy, benzyloxy, NH-phenyl,  $NH-C_1-C_2alkyl-C(=O)-O-C_1-C_2alkyl$ -phenyl,  $-P(=O)(OC_1-C_6alkyl)_2$ ; or phenyl, phenoxy, benzyloxy or NH-phenyl substituted by chlorine, fluorine, methoxy, trifluoromethyl or trifluoromethoxy;

more especially wherein  $R_5$  is  $C_1$ - $C_{12}alkoxy$ ;

(19) compounds according to one of groups (1) to (6) of formula (I) wherein  $R_3$  is  $-C_2-C_6alkyl-NHC(=O)R_6$  and  $R_6$  is H,  $C_1$ - $C_4alkyl$ ,  $C_2$ - $C_4alkenyl$ ,  $C_2$ - $C_4alkynyl$ , phenyl or benzyl;

(20) compounds according to one of groups (1) to (6) of formula (I) wherein  $R_3$  is  $-CH_2$ -heterocyclyl, and heterocyclyl denotes pyridyl, furyl, tetrahydrofuranyl, pyranal, tetrahydropyranal, pyrazolyl, imidazolyl, thiazolyl, benzothienyl, quinoliny, quinoxaliny, benzofuranyl, benzimidazolyl, benzopyrrolyl, benzothiazolyl, indolyl, coumariny or indazolyl, the mentioned radicals being unsubstituted or substituted by one or two substituents selected independently of one another from halogen, trifluoromethyl, trifluoromethoxy and nitro; especially pyridyl, furyl, pyrazolyl, imidazolyl, thiazolyl, benzimidazolyl, benzopyrrolyl, benzothiazolyl or indolyl unsubstituted or substituted by one or two substituents selected independently of one another from halogen, trifluoromethyl, trifluoromethoxy and nitro; more especially pyridyl or thiazolyl unsubstituted or substituted by one or two substituents selected independently of one another from halogen, trifluoromethyl, trifluoromethoxy and nitro, especially mono-substituted by chlorine;

(21) compounds according to one of groups (1) to (6) of formula (I) wherein  $R_3$  is  $C_2$ - $C_{10}alkenyl$ , especially  $C_2$ - $C_4alkenyl$ , unsubstituted or mono- or di-substituted, especially mono-substituted, by  $C_2$ - $C_4alkynyl$ ,  $-C(=O)-C_1-C_4alkoxy$ ,  $-C(=O)-O-C_1-C_4alkyl$ -benzoyl, phenyl or halogen; more especially wherein  $R_3$  is  $-CH_2-CH=CH_2$ ;

(22) compounds according to one of groups (1) to (6) of formula (I) wherein  $R_3$  is branched, unsubstituted  $C_4$ - $C_{10}$ alkyl;

(23) compounds according to one of groups (1) to (6) of formula (I) wherein  $R_3$  is branched, substituted  $C_3$ - $C_{10}$ alkyl;

(24) compounds according to one of groups (1) to (6) of formula (I) wherein  $R_3$  is unsubstituted benzyl;

(25) compounds according to one of groups (1) to (24) of formula (I) wherein  $X^{n-}$  is benzoate;

(26) compounds according to one of groups (1) to (24) of formula (I) wherein

$X^{n-}$  is a substituted benzoate, especially a mono- or di-substituted benzoate; especially wherein the benzoate is substituted by one or two substituents selected independently of one another from  $C_1$ - $C_4$ alkyl, halo- $C_1$ - $C_4$ alkyl, OH, SH, halogen and phenyl; more especially mono-substituted by methyl, tert-butyl, trifluoromethyl, OH, SH, fluorine or phenyl;

(27) compounds according to one of groups (1) to (24) of formula (I) wherein

$X^{n-}$  is the anion of a sulfonic acid; especially a halo-substituted  $C_1$ - $C_4$ alkane- or aryl-sulfonic acid; more especially trifluoromethylsulfonic acid or benzenesulfonic acid;

(28) compounds according to one of groups (1) to (24) of formula (I) wherein

$X^{n-}$  is the anion of a  $C_1$ - $C_4$ alkane- or  $C_1$ - $C_4$ alkene-monocarboxylic acid unsubstituted or substituted by OH,  $C_1$ - $C_4$ alkoxy or phenoxy;

(29) compounds according to one of groups (1) to (24) of formula (I) wherein

$X^{n-}$  is the anion of an unsubstituted or substituted dicarboxylic acid; especially tartaric acid, maleic acid or 2,2'-oxydiacetic acid;

(30) compounds according to one of groups (1) to (24) of formula (I) wherein

$X^{n-}$  is the anion of an unsubstituted or substituted tricarboxylic acid, especially citric acid;

(31) compounds according to one of groups (1) to (24) of formula (I) wherein

$X^{n-}$  is the anion of an acid of the formula Het-COOH wherein Het is a heterocyclic ring; especially wherein Het is furyl;

(32) compounds according to one of groups (1) to (24) of formula (I) wherein

$X^{n-}$  is the anion of an inorganic acid, especially sulfate or hydrogen sulfate, more especially sulfate;


(33) compounds according to one of groups (1) to (24), (29) or (30) of formula (I) wherein

n is 1 or 2, especially 1.

Special preference within the context of the invention is given to the compounds of formula (I) listed in the Tables and, where applicable, their E/Z isomers and mixtures of E/Z isomers; more especially salts of the compounds

4"-deoxy-4"-epi-amino-avermectin B1;  
4"-deoxy-4"-epi-dimethyl-amino-avermectin B1;  
4"-deoxy-4"-epi-N-ethylamino-avermectin B1;  
4"-deoxy-4"-epi-N-prop-1-ylamino-avermectin B1;  
4"-deoxy-4"-epi-(N-ethyl-N-methyl-amino)-avermectin B1;  
4"-deoxy-4"-epi-(N-methyl-N-prop-1-yl-amino)-avermectin B1;  
4"-deoxy-4"-epi-(N-isopropyl-N-methylamino)-avermectin B1;  
4"-deoxy-4"-epi-(N-methyl-N-1-propen-3-yl-amino)-avermectin B1;  
4"-deoxy-4"-epi-(N-methyl-N-ethoxycarbonylmethyl-amino)-avermectin B1;  
4"-deoxy-4"-epi-(N-methyl-N-benzyl-amino)-avermectin B1;  
4"-deoxy-4"-epi-(N-methyl-N-4-difluoromethoxyphenylmethyl-amino)-avermectin B1;  
4"-deoxy-4"-epi-(N-methyl-N-2,5-dichlorophenylmethyl-amino)-avermectin B1;  
4"-deoxy-4"-epi-(N-methyl-N-2,5-difluorophenylmethyl-amino)-avermectin B1;  
4"-deoxy-4"-epi-(N-methyl-N-2,3,4-trifluorophenylmethyl-amino)-avermectin B1;  
4"-deoxy-4"-epi-(pyrrolidin-1-yl)-avermectin B1; and  
4"-deoxy-4"-epi-(azetidin-1-yl)-avermectin B1.

The invention relates also to a process for the preparation of the compounds of formula (I) and, where applicable, their tautomers, which process comprises first of all preparing a compound of formula



wherein R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are as defined for formula (I) above; for example as follows:

(A) for the preparation of a compound of formula (Ia) wherein R<sub>1</sub> is as defined above under (1) for formula (I), R<sub>2</sub> is hydrogen and R<sub>3</sub> is a group R<sub>31</sub>-CH-R<sub>32</sub>, wherein R<sub>31</sub> is C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl, heterocyclyl or unsubstituted or substituted C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl or heterocyclyl, and R<sub>32</sub> is H or unsubstituted or substituted C<sub>1</sub>-C<sub>5</sub>alkyl;

a compound of formula (Ia) in which R<sub>1</sub> is as defined above under (1) for formula (I) and R<sub>2</sub> and R<sub>3</sub> are hydrogen and which can be prepared according to methods known *per se* is reacted in the presence of a reducing agent with a compound R<sub>31</sub>-C(=O)-R<sub>32</sub>, wherein R<sub>31</sub> and R<sub>32</sub> are as defined above; or

(B) for the preparation of a compound of formula (Ia) wherein R<sub>1</sub> and R<sub>2</sub> are as defined above under (1) for formula (I), and R<sub>3</sub> is as defined above under (1) for formula (I) with the exception of hydrogen, a compound of formula (Ia) in which R<sub>1</sub> and R<sub>2</sub> are as defined above under (1) for formula (I) and R<sub>3</sub> is hydrogen and which can be prepared according to methods known *per se*,

is reacted with a compound of formula R<sub>3</sub>-Hal wherein R<sub>3</sub> is as defined above under (1) for formula (I) and Hal is halogen, especially bromine or iodine; or

(C) for the preparation of a compound of formula (Ia) wherein R<sub>1</sub> and R<sub>2</sub> are as defined above under (1) for formula (I) and R<sub>3</sub> is hydroxy-substituted -CH<sub>2</sub>-C<sub>1</sub>-C<sub>11</sub>alkyl,

a compound of formula (Ia) wherein R<sub>1</sub> and R<sub>2</sub> are as defined above under (1) for formula (I), R<sub>3</sub> is -C(=O)-R<sub>5</sub>-substituted -C<sub>1</sub>-C<sub>11</sub>alkyl and R<sub>5</sub> is OH or alkoxy is reacted with a reducing agent; or

(D) for the preparation of a compound of formula (Ia) wherein  $R_1$  and  $R_2$  are as defined above under (1) for formula (I) and  $R_3$  is COOH-substituted  $C_1$ - $C_{12}$ alkyl,

a compound of formula (Ia) wherein  $R_1$  and  $R_2$  are as defined above under (1) for formula (I),  $R_3$  is  $-C(=O)-R_5$ -substituted  $C_1$ - $C_{12}$ alkyl and  $R_5$  is  $C_1$ - $C_6$ alkoxy or benzyloxy is reacted with a base or with a reducing agent; or

(E) for the preparation of a compound of formula (Ia) wherein  $R_1$  and  $R_3$  are as defined above under (1) for formula (I) and  $R_2$  is methyl,

a compound of formula (Ia) wherein  $R_1$  and  $R_3$  are as defined above under (1) for formula (I) and  $R_2$  is hydrogen is reacted with a compound of the formula methyl-Hal, wherein Hal is a halogen; or with formaldehyde in the presence of a reducing agent; or

(F) for the preparation of a compound of formula (Ia) wherein  $R_1$  and  $R_2$  are as defined above under (1) for formula (I) and  $R_3$  is  $-C(=O)N(R_8)_2$ -substituted  $C_1$ - $C_{12}$ alkyl and wherein the two  $R_8$  are each independently of the other H or unsubstituted or substituted  $C_1$ - $C_{12}$ alkyl,

a compound of formula (Ia) wherein  $R_1$  and  $R_2$  are as defined above under (1) for formula (I) and  $R_3$  is  $-C(=O)R_5$ -substituted  $C_1$ - $C_{12}$ alkyl and  $R_5$  is OH is reacted with a compound of formula  $NH(R_8)_2$  wherein  $R_8$  is H or unsubstituted or substituted  $C_1$ - $C_{12}$ alkyl, in the presence of a water-removing agent; or

(G) for the preparation of a compound of formula (Ia) wherein  $R_1$  and  $R_2$  are as defined above under (1) for formula (I) and  $R_3$  is hydroxy-substituted  $C_4$ - $C_{12}$ alkyl,

a compound of formula (Ia) wherein  $R_1$  and  $R_2$  are as defined above under (1) for formula (I),  $R_3$  is  $-C(=O)-R_5$ -substituted  $C_1$ - $C_5$ alkyl and  $R_5$  is  $C_1$ - $C_{12}$ alkoxy is reacted with two moles of a  $C_1$ - $C_3$ alkylmagnesium halide or  $C_1$ - $C_3$ alkyllithium reagent; or

(H) for the preparation of a compound of formula (Ia) wherein  $R_1$  is as defined above under (1) for formula (I) and  $R_2$  and  $R_3$  together are a three- to seven-membered alkylene or four- to seven-membered alkenylene bridge, and wherein a  $CH_2$  group may have been replaced by O, S or  $NR_4$ , and  $R_4$  is as defined above under (1) for formula (I),

a compound of formula (Ia) wherein  $R_1$  is as defined above under (1) for formula (I) and  $R_2$  and  $R_3$  are hydrogen is reacted with a compound of formula Hal-( $C_3$ - $C_7$ alkylene)-Hal or Hal-( $C_4$ - $C_7$ alkenylene)-Hal, wherein Hal is a halogen, and wherein a  $CH_2$  group may have been replaced by O, S or  $NR_4$ , and  $R_4$  is as defined above under (1) for formula (I); or

(I) for the preparation of a compound of formula (Ia) wherein  $R_1$  is as defined above under (1) for formula (I) and  $R_2$  and  $R_3$  are identical and are as defined under (1) for formula (I),

a compound of formula (Ia) wherein  $R_1$  is as defined under (1) for formula (I) and  $R_2$  and  $R_3$  are hydrogen is reacted with two moles of a compound of formula  $R_3$ -Hal wherein  $R_3$  is as defined above for formula (I) and Hal is halogen, preferably bromine or iodine; or

(J) for the preparation of a compound of formula (Ia) wherein  $R_2$  and  $R_3$  are identical and are unsubstituted or mono- to penta-substituted  $-\text{CH}_2\text{-C}_1\text{-C}_{11}\text{alkyl}$ , unsubstituted or mono- to penta-substituted  $-\text{CH}_2\text{-C}_1\text{-C}_{11}\text{alkenyl}$  or unsubstituted or mono- to penta-substituted  $-\text{CH}_2\text{-C}_1\text{-C}_{11}\text{alkynyl}$ ,

a compound of formula (Ia) wherein  $R_1$  is as defined above under (1) for formula (I) and  $R_2$  and  $R_3$  are hydrogen is reacted with two moles of a compound of formula  $R_{31}\text{-CHO}$  wherein  $R_{31}$  is unsubstituted or mono- to penta-substituted  $\text{C}_1\text{-C}_{11}\text{alkyl}$ , unsubstituted or mono- to penta-substituted  $\text{C}_1\text{-C}_{11}\text{alkenyl}$  or unsubstituted or mono- to penta-substituted  $\text{C}_1\text{-C}_{11}\text{alkynyl}$ , in the presence of a reducing agent; and then

(K) a compound of formula (Ia) prepared, for example, in accordance with any one of processes A) to J) mentioned above is reacted with an acid  $\text{XH}_n$  wherein X and n are as defined above under formula (I).

The comments made above in connection with tautomers of compounds of formula (I) apply analogously to the starting materials mentioned hereinabove and hereinbelow in respect of their tautomers.

The reactions described hereinabove and hereinbelow are carried out in a manner known *per se*, for example in the absence or, customarily, in the presence of a suitable solvent or diluent or of a mixture thereof, the reactions being carried out, as required, with cooling, at room temperature or with heating, for example in a temperature range of approximately from  $-80^\circ\text{C}$  to the boiling temperature of the reaction medium, preferably from approximately  $0^\circ\text{C}$  to approximately  $+150^\circ\text{C}$ , and, if necessary, in a closed vessel, under pressure, under an inert gas atmosphere and/or under anhydrous conditions. Especially advantageous reaction conditions can be found in the Examples.

The reaction time is not critical; a reaction time of from about 0.1 to about 24 hours, especially from about 0.5 to about 10 hours, is preferred.

The product is isolated by customary methods, for example by means of filtration, crystallisation, distillation or chromatography, or any suitable combination of such methods.

The starting materials mentioned hereinabove and hereinbelow that are used for the preparation of the compounds of formula (I) and, where applicable, their tautomers are known or can be prepared by methods known *per se*, e.g. as indicated below.

Process variant (A):

Examples of solvents and diluents include: aromatic, aliphatic and alicyclic hydrocarbons and halogenated hydrocarbons, such as benzene, toluene, xylene, mesitylene, Tetralin, chlorobenzene, dichlorobenzene, bromobenzene, petroleum ether, hexane, cyclohexane, dichloromethane, trichloromethane, tetrachloromethane, dichloroethane, trichloroethene or tetrachloroethene; ethers, such as diethyl ether, dipropyl ether, diisopropyl ether, dibutyl ether, tert-butyl methyl ether, ethylene glycol monomethyl ether, ethylene glycol monoethyl ether, ethylene glycol dimethyl ether, dimethoxydiethyl ether, tetrahydrofuran or dioxane; alcohols, such as methanol, ethanol, propanol, isopropanol, butanol, ethylene glycol or glycerol; carboxylic acid, such as acetic acid or formic acid; amides, such as N,N-dimethylformamide, N,N-diethylformamide, N,N-dimethylacetamide, N-methylpyrrolidone or hexamethylphosphoric acid triamide; nitriles, such as acetonitrile or propionitrile; and sulfoxides, such as dimethyl sulfoxide; and also water; or mixtures of the mentioned solvents;

especially suitable are ethers, alcohols, water and carboxylic acids, more especially tetrahydrofuran, acetic acid or water.

The reactions are advantageously carried out in a temperature range of from about room temperature to the boiling point of the solvent used; preference being given to reaction at 10 to 30°C.

In a preferred embodiment of Variant (A) the reaction is carried out at room temperature, in tetrahydrofuran in the presence of acetic acid. Especially preferred conditions for the reaction are described in Example P1.1.

Process variant (B):

Examples of solvents and diluents include: aromatic, aliphatic and alicyclic hydrocarbons and halogenated hydrocarbons and ethers as listed above under Process variant (A); ketones, such as acetone, methyl ethyl ketone or methyl isobutyl ketone; alcohols, such as methanol, ethanol, propanol, isopropanol, butanol, ethylene glycol or

glycerol; carboxylic acid esters, such as methyl acetate, ethyl acetate, or esters of benzoic acid; amides as listed above under Process variant (A); nitriles, such as acetonitrile or propionitrile; and sulfoxides, such as dimethyl sulfoxide; and also water; or mixtures of the mentioned solvents;

especially suitable are water, esters of organic acids, halogenated hydrocarbons and aromatic hydrocarbons; more especially two-phase mixtures of such an organic solvent with water.

The reactions are advantageously carried out in a temperature range of approximately from room temperature to the boiling point of the solvent used, preferably from room temperature up to 90°C, especially up to 60°C, and in the presence of a base, preferably an inorganic base, for example sodium hydroxide, potassium hydroxide, sodium carbonate or sodium hydrogen carbonate.

Especially preferred conditions for the reaction are described, for example, in Examples P1.2, P1.3, P2.1 and P2.7.

Process variant (C):

Examples of solvents and diluents include: aromatic, aliphatic and alicyclic hydrocarbons and halogenated hydrocarbons and ethers, amides and nitriles as listed above under Process variant (A); and sulfoxides, such as dimethyl sulfoxide; or mixtures of the mentioned solvents; ethers and hydrocarbons being especially suitable.

The reactions are advantageously carried out in a temperature range of from 0°C to the boiling point of the solvent used, preferably from 0°C to room temperature. Especially preferred conditions for the reaction are described, for example, in Example P2.2.

Process variant (D):

Suitable solvents include those mentioned under Variant (A); additionally also ketones, such as acetone, methyl ethyl ketone and methyl isobutyl ketone; and carboxylic acids, such as acetic acid or formic acid; carboxylic acid esters, such as methyl acetate, ethyl acetate, or esters of benzoic acid.

The reactions are advantageously carried out in a temperature range of approximately from room temperature to the boiling point of the solvent used, preferably in the presence of an inorganic base, for example lithium hydroxide, sodium hydroxide, potassium hydroxide, sodium carbonate or sodium hydrogen carbonate.



Especially preferred conditions for this Process variant are described, for example, in Example P2.6.

As an alternative it is possible to choose a reaction variant wherein a reducing agent, especially molecular hydrogen, is used, more especially in a mixture of tetrahydrofuran and water as solvent and in the presence of a heavy metal catalyst, especially a Pd catalyst.

Especially preferred conditions for this Process variant are described, for example, in Example P2.5.

Process variant (E):

Suitable solvents include those mentioned under Variant (B), especially suitable solvents being esters of organic acids, halogenated hydrocarbons and aromatic hydrocarbons; especially two-phase mixtures of an ester with water.

The reactions are advantageously carried out in a temperature range of from 0°C to the boiling point of the solvent used, preferably from room temperature to 60°C, and in the presence of a base, preferably an inorganic base, for example sodium hydroxide, potassium hydroxide, sodium carbonate or sodium hydrogen carbonate.

Especially preferred conditions for this Process variant are described, for example, in Example P2.3.

In an alternative embodiment, suitable solvents include those mentioned above, preferably ethers, alcohols, water and carboxylic acids, in combination with a hydride, such as a borohydride, especially  $\text{NaCNBH}_3$ .

Especially preferred conditions for this Process variant are described, for example, in Example P2.4.

Process variant (F):

Examples of solvents and diluents include: aromatic, aliphatic and alicyclic hydrocarbons and halogenated hydrocarbons; ethers, amides and nitriles as listed above under Process variant (A); ketones, such as acetone, methyl ethyl ketone or methyl isobutyl ketone; carboxylic acid esters, such as methyl acetate, ethyl acetate, or esters of benzoic acid; and sulfoxides, such as dimethyl sulfoxide; or mixtures of the mentioned solvents;

especially suitable are esters of organic acids, such as ethyl acetate.

As water-removing agent there are used the customary peptide coupling reagents, especially carbodiimides and hydroxybenzotriazoles.

The reactions are advantageously carried out in a temperature range of from 0°C to the boiling point of the solvent used, preferably at room temperature.

Especially preferred conditions for the reaction are described, for example, in Example P2.8.

Process variant (G):

Examples of solvents and diluents include: aromatic, aliphatic and alicyclic hydrocarbons and ethers as listed above under Process variant (A); and sulfoxides, such as dimethyl sulfoxide; or mixtures of the mentioned solvents; ethers, more especially tetrahydrofuran, being especially suitable.

The reactions are advantageously carried out in a temperature range of from 0°C to the boiling point of the solvent used, preferably from 0°C to room temperature.

Especially preferred conditions for the reaction are described, for example, in Example P2.10.

Process variant (H):

Suitable solvents include those mentioned under Variant (B), especially suitable solvents being water, esters of organic acids, halogenated hydrocarbons and aromatic hydrocarbons; especially two-phase mixtures of such an organic solvent with water.

The reactions are advantageously carried out in a temperature range of from 0°C to the boiling point of the solvent used, preferably from 90°C to the boiling point of the solvent, and in the presence of a base, preferably an inorganic base, for example sodium hydroxide, potassium hydroxide, sodium carbonate or sodium hydrogen carbonate.

Especially preferred conditions for the reaction are described, for example, in Example P3.2.

Process variant (I):

Suitable solvents include those mentioned under Variant (B), especially suitable solvents being water, esters of organic acids, halogenated hydrocarbons and aromatic hydrocarbons; especially two-phase mixtures of such an organic solvent with water.

The reactions are advantageously carried out in a temperature range of from 0°C to the boiling point of the solvent used, preferably from 90°C to the boiling point, and in the presence of a base, preferably an inorganic base, for example sodium hydroxide, potassium hydroxide, sodium carbonate or sodium hydrogen carbonate.

Especially preferred conditions for the reaction are described, for example, in Example P3.1.

Process variant (J):

Suitable solvents include those mentioned under Variant (B), especially suitable solvents being water, ethers of organic acids, alcohols and water; especially two-phase mixtures of an ether with water.

The reactions are advantageously carried out in a temperature range of from 0°C to the boiling point of the solvent used, preferably at room temperature.

Especially preferred conditions for the reaction are described, for example, in Example P3.3.

Process variant (K):

Especially suitable solvents are listed under Process variant (B); dichloromethane, acetonitrile, ethyl acetate, toluene and dioxane being especially suitable.

The operation is preferably carried out in a temperature range of from 0°C to the boiling point of the solvent, preferably at from 0°C to room temperature.

Especially preferred conditions for the reaction are described in Examples P4.1 to P5.5.

The compounds of formula (I) may be in the form of one of the possible isomers or in the form of a mixture thereof, in the form of pure isomers or in the form of an isomeric mixture, i.e. in the form of a racemic mixture; the invention relates both to the pure isomers and to the racemic mixtures and is to be interpreted accordingly hereinabove and hereinbelow, even if stereochemical details are not mentioned specifically in every case.

The racemates can be resolved into the optical antipodes by known methods, for example by recrystallisation from an optically active solvent, by chromatography on chiral adsorbents, for example high pressure liquid chromatography (HPLC) on acetylcellulose, with the aid of suitable microorganisms, by cleavage with specific, immobilised enzymes, or *via* the formation of inclusion compounds, for example using chiral crown ethers, only one isomer being complexed.

Apart from by separation of corresponding mixtures of isomers, pure optical isomers can be obtained according to the invention also by generally known methods of enantio-

selective synthesis, for example by carrying out the process according to the invention using starting materials having correspondingly suitable stereochemistry.

In each case it is advantageous to isolate or synthesise the biologically more active isomer, where the individual components have different biological activity.

The compounds of formula (I) may also be obtained in the form of their hydrates and/or may include other solvents, for example solvents which may have been used for the crystallisation of compounds in solid form.

The invention relates to all those embodiments of the process according to which a compound obtainable as starting material or intermediate at any stage of the process is used as starting material and some or all of the remaining steps are carried out or a starting material is used in the form of a derivative or salt and/or its racemates or antipodes or, especially, is formed under the reaction conditions.

In the processes of the present invention it is preferable to use those starting materials and intermediates which result in the compounds of formula (I) that are especially preferred.

The invention relates especially to the preparation processes described in Examples P1.1 to P5.5.

In the area of pest control, the compounds of formula (I) according to the invention are active ingredients exhibiting valuable preventive and/or curative activity with a very advantageous biocidal spectrum and a very broad spectrum, even at low rates of concentration, while being well tolerated by warm-blooded animals, fish and plants. They are, surprisingly, equally suitable for controlling both plant pests and ecto- and endo-parasites in humans and more especially in productive livestock, domestic animals and pets. They are effective against all or individual development stages of normally sensitive animal pests, but also of resistant animal pests, such as insects and representatives of the order Acarina, nematodes, cestodes and trematodes, while at the same time protecting useful organisms. The insecticidal or acaricidal activity of the active ingredients according to the invention may manifest itself directly, i.e. in the mortality of the pests, which occurs immediately or only after some time, for example during moulting, or indirectly, for example in reduced oviposition and/or hatching rate, good activity corresponding to a mortality of at least 50 to 60 %.

The action of the compounds according to the invention and the compositions comprising them against animal pests can be significantly broadened and adapted to the

given circumstances by the addition of other insecticides, acaricides or nematicides. Suitable additives include, for example, representatives of the following classes of active ingredient: organophosphorus compounds, nitrophenols and derivatives, formamidines, ureas, carbamates, pyrethroids, chlorinated hydrocarbons and *Bacillus thuringiensis* preparations.

Examples of especially suitable mixing partners include: azamethiphos; chlorfenvinphos; bupirimate; cypermethrin, cypermethrin high-cis; cyromazine; diafenthiuron; diazinon; dichlorvos; dicrotophos; dicyclanil; fenoxycarb; fluazuron; furathiocarb; isazofos; iodfenphos; kinoprene; lufenuron; methacriphos; methidathion; monocrotophos; phosphamidon; profenofos; diofenolan; a substance obtainable from the *Bacillus thuringiensis* strain GC91 or from NCTC11821; pymetrozine; bromopropylate; methoprene; disulfoton; quinalphos; tau-fluvalinate; thiocyclam; thiometon; aldicarb; azinphos-methyl; benfuracarb; bifenthrin; buprofezin; carbofuran; dibutylaminothio; cartap; chlorfluazuron; chlorpyrifos; cyfluthrin; alpha-cypermethrin; zeta-cypermethrin; deltamethrin; diflubenzuron; endosulfan; ethiofencarb; fenitrothion; fenazaquin; fenobucarb; fenvalerate; formothion; methiocarb; heptenophos; imidacloprid; isoprocarb; methamidophos; methomyl; mevinphos; parathion; parathion-methyl; phosalone; pirimicarb; propoxur; teflubenzuron; terbufos; triazamate; abamectin; fenobucarb; tebufenozide; fipronil; beta-cyfluthrin; silafluofen; fenpyroximate; pyridaben; primicarb; pyriproxyfen; pyrimidifen; nematorin; nitenpyram; NI-25, acetamiprid; avermectin B<sub>1</sub> (abamectin); an insect-active extract from a plant; a preparation comprising insect-active nematodes; a preparation obtainable from *Bacillus subtilis*; a preparation comprising insect-active fungi; a preparation comprising insect-active viruses; AC 303 630; acephate; acrinathrin; alanycarb; alphamethrin; amitraz; AZ 60541; azinphos A; azinphos M; azocyclotin; bendiocarb; bensultap; betacyfluthrin; BPMC; brofenprox; bromophos A; bufencarb; butocarboxim; butylpyridaben; cadusafos; carbaryl; carbophenothion; chloethiocarb; chlorethoxyfos; chlormephos; cis-res-methrin; clocythrins; clofentezine; cyanophos; cycloprothrin; cyhexatin; demeton M; demeton S; demeton-S-methyl; dichlofenthion; dicliphos; diethion; dimethoate; dimethylvinphos; dioxathion; edifenphos; emamectin; esfenvalerate; ethion; ethofenprox; ethoprophos; etrimphos; fenamiphos; fenbutatin oxide; fenothiocarb; fenpropathrin; fenpyrad; fenthion; fluazinam; flucycloxuron; flucythrinate; flufenoxuron; flufenprox; fonophos; fosthiazate; fubfenprox; HCH; hexaflumuron; hexythiazox; IKI-220; iprobenfos; isofenphos; isoxathion; ivermectin; lambda-cyhalothrin; malathion; mecarbam; mesulfenphos; metaldehyd; metolcarb; milbemectin; moxidectin; naled; NC 184; omethoate; oxamyl; oxydemethon M; oxydeprofos; permethrin; phenthoate; phorate; phosmet; phoxim; pirimiphos M; pirimiphos A; promecarb; propaphos; prothiofos;

prothoate; pyrachlophos; pyrada-phenthion; pyresmethrin; pyrethrum; RH 5992; salithion; sebufos; sulfotep; sulprofos; tebufenpyrad; tebupirimphos; tefluthrin; temephos; terbam; tetrachlorvinphos; thiacloprid; thiamethoxam; thiafenox; thiodicarb; thiofanox; thionazin; thuringiensin; tralomethrin; triarathen; triazophos; triazuron; trichlorfon; triflumuron; trimethacarb; vamidothion; xylylcarb; YI 5301/5302; zetamethrin; DPX-MP062; RH-2485; D 2341 or XMC (3,5-xylyl methylcarbamate).

The said animal pests include, for example, those mentioned in European Patent Application EP-A-736 252, page 5, line 55, to page 6, line 55. The pests mentioned therein are therefore included by reference in the subject matter of the present invention.

It is also possible to control pests of the class Nematoda using the compounds according to the invention. Such pests include, for example,

root knot nematodes, cyst-forming nematodes and also stem and leaf nematodes; especially of *Heterodera* spp., e.g. *Heterodera schachtii*, *Heterodora avenae* and *Heterodora trifolii*; *Globodera* spp., e.g. *Globodera rostochiensis*; *Meloidogyne* spp., e.g. *Meloidogyne incognita* and *Meloidogyne javanica*; *Radopholus* spp., e.g. *Radopholus simiis*; *Pratylenchus*, e.g. *Pratylenchus neglectans* and *Pratylenchus penetrans*; *Tylenchulus*, e.g. *Tylenchulus semipenetrans*; *Longidorus*, *Trichodorus*, *Xiphinema*, *Ditylenchus*, *Aphelenchoides* and *Anguina*; insbesondere *Meloidogyne*, e.g. *Meloidogyne incognita*, and *Heterodera*, e.g. *Heterodera glycines*.

An especially important aspect of the present invention is the use of the compounds of formula (I) according to the invention in the protection of plants against parasitic feeding pests.

The compounds according to the invention can be used to control, i.e. to inhibit or destroy, pests of the mentioned type occurring on plants, especially on useful plants and ornamentals in agriculture, in horticulture and in forestry, or on parts of such plants, such as the fruits, blossoms, leaves, stems, tubers or roots, while in some cases plant parts that grow later are still protected against those pests.

Target crops include especially cereals, such as wheat, barley, rye, oats, rice, maize and sorghum; beet, such as sugar beet and fodder beet; fruit, e.g. pomes, stone fruit and soft fruit, such as apples, pears, plums, peaches, almonds, cherries and berries, e.g. strawberries, raspberries and blackberries; leguminous plants, such as beans, lentils, peas and soybeans; oil plants, such as rape, mustard, poppy, olives, sunflowers, coconut, castor oil,

cocoa and groundnuts; cucurbitaceae, such as marrows, cucumbers and melons; fibre plants, such as cotton, flax, hemp and jute; citrus fruits, such as oranges, lemons, grapefruit and mandarins; vegetables, such as spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes and paprika; lauraceae, such as avocado, cinnamon and camphor; and tobacco, nuts, coffee, aubergines, sugar cane, tea, pepper, vines, hops, bananas, natural rubber plants and ornamentals.

Further areas of use of the compounds according to the invention are the protection of stored goods and storerooms and the protection of raw materials, and also in the hygiene sector, especially the protection of domestic animals and productive livestock against pests of the mentioned type, more especially the protection of domestic animals, especially cats and dogs, from attack by fleas, ticks and nematodes.

The invention therefore relates also to pesticidal compositions, such as emulsifiable concentrates, suspension concentrates, directly sprayable or dilutable solutions, spreadable pastes, dilute emulsions, wettable powders, soluble powders, dispersible powders, wettable powders, dusts, granules and encapsulations of polymer substances, that comprise at least one of the compounds according to the invention, the choice of formulation being made in accordance with the intended objectives and the prevailing circumstances.

The active ingredient is used in those compositions in pure form, a solid active ingredient, for example, in a specific particle size, or preferably together with at least one of the adjuvants customary in formulation technology, such as extenders, e.g. solvents or solid carriers, or surface-active compounds (surfactants). In the area of parasite control in humans, domestic animals, productive livestock and pets it will be self-evident that only physiologically tolerable additives are used.

As formulation adjuvants there are used, for example, solid carriers, solvents, stabilisers, "slow release" adjuvants, colourings and optionally surface-active substances (surfactants). Suitable carriers and adjuvants include all substances customarily used. As adjuvants, such as solvents, solid carriers, surface-active compounds, non-ionic surfactants, cationic surfactants, anionic surfactants and further adjuvants in the compositions used according to the invention, there come into consideration, for example, those described in EP-A-736 252, page 7, line 51 to page 8, line 39.

The compositions for use in crop protection and in humans, domestic animals and productive livestock generally comprise from 0.1 to 99 %, especially from 0.1 to 95 %, of active ingredient and from 1 to 99.9 %, especially from 5 to 99.9 %, of at least one solid or

liquid adjuvant, the composition generally including from 0 to 25 %, especially from 0.1 to 20 %, of surfactants (% = % by weight in each case). Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations having considerably lower concentrations of active ingredient.

Preferred crop protection products have especially the following compositions (% = percent by weight):

Emulsifiable concentrates:

active ingredient:	1 to 90%, preferably 5 to 20%
surfactant:	1 to 30%, preferably 10 to 20%
solvent:	5 to 98%, preferably 70 to 85%

Dusts:

active ingredient:	0.1 to 10%, preferably 0.1 to 1%
solid carrier:	99.9 to 90%, preferably 99.9 to 99%

Suspension concentrates:

active ingredient:	5 to 75%, preferably 10 to 50%
water:	94 to 24%, preferably 88 to 30%
surfactant:	1 to 40%, preferably 2 to 30%

Wettable powders:

active ingredient:	0.5 to 90%, preferably 1 to 80%
surfactant:	0.5 to 20%, preferably 1 to 15%
solid carrier:	5 to 99%, preferably 15 to 98%

Granules:

active ingredient:	0.5 to 30%, preferably 3 to 15%
solid carrier:	99.5 to 70%, preferably 97 to 85%

The compositions according to the invention may also comprise further solid or liquid adjuvants, such as stabilisers, e.g. vegetable oils or epoxidised vegetable oils (e.g. epoxidised coconut oil, rapeseed oil or soybean oil), antifoams, e.g. silicone oil, preservatives, viscosity regulators, binders and/or tackifiers as well as fertilisers or other active ingredients for obtaining special effects, e.g. acaricides, bactericides, fungicides, nematocides, molluscicides or selective herbicides.



The crop protection products according to the invention are prepared in known manner, in the absence of adjuvants, e.g. by grinding, sieving and/or compressing a solid active ingredient or mixture of active ingredients, for example to a certain particle size, and in the presence of at least one adjuvant, for example by intimately mixing and/or grinding the active ingredient or mixture of active ingredients with the adjuvant(s). The invention relates likewise to those processes for the preparation of the compositions according to the invention and to the use of the compounds of formula (I) in the preparation of those compositions.

The invention relates also to the methods of application of the crop protection products, i.e. the methods of controlling pests of the mentioned type, such as spraying, atomising, dusting, coating, dressing, scattering or pouring, which are selected in accordance with the intended objectives and the prevailing circumstances, and to the use of the compositions for controlling pests of the mentioned type. Typical rates of concentration are from 0.1 to 1000 ppm, preferably from 0.1 to 500 ppm, of active ingredient. The rates of application per hectare are generally from 1 to 2000 g of active ingredient per hectare, especially from 10 to 1000 g/ha, preferably from 20 to 600 g/ha.

A preferred method of application in the area of crop protection is application to the foliage of the plants (foliar application), the frequency and the rate of application being dependent upon the risk of infestation by the pest in question. However, the active ingredient can also penetrate the plants through the roots (systemic action) when the locus of the plants is impregnated with a liquid formulation or when the active ingredient is incorporated in solid form into the locus of the plants, for example into the soil, e.g. in granular form (soil application). In the case of paddy rice crops, such granules may be applied in metered amounts to the flooded rice field.

The crop protection products according to the invention are also suitable for protecting plant propagation material, e.g. seed, such as fruits, tubers or grains, or plant cuttings, against animal pests. The propagation material can be treated with the composition before planting: seed, for example, can be dressed before being sown. The active ingredients according to the invention can also be applied to grains (coating), either by impregnating the seeds in a liquid formulation or by coating them with a solid formulation. The composition can also be applied to the planting site when the propagation material is being planted, for example to the seed furrow during sowing. The invention relates also to such methods of treating plant propagation material and to the plant propagation material so treated.

Preparation Examples: Preparation of the free basesExample P1.1: 4"-Desoxy-4"-epi-(N-3-fluorophenylmethyl-amino)-avermectin B1

1.0 g of 4"-desoxy-4"-epi-amino-avermectin B1 is dissolved in 12 ml of tetrahydrofuran. 1.8 ml of acetic acid, 0.2 ml of water and 0.18 ml of 3-fluorobenzaldehyde are added. 90 mg of sodium cyanoborohydride are then added. The mixture is stirred at room temperature for 12 hours. Extraction is then carried out with ethyl acetate and saturated sodium chloride solution; the organic phase is dried over sodium sulfate and the solvents are distilled off. The residue is purified by chromatography on silica gel (hexane/ethyl acetate), yielding 4"-desoxy-4"-epi-(N-3-fluorophenylmethyl-amino)-avermectin B1.

Example P1.2: 4"-Desoxy-4"-epi-N-ethylamino-avermectin B1

4.0 g of 4"-desoxy-4"-epi-amino-avermectin B1 are dissolved in 24 ml of ethyl acetate. 7.4 ml of ethyl iodide and 24 ml of sodium bicarbonate (1N in water) are added. The mixture is stirred vigorously at room temperature for 14 hours. The phases are then separated; the organic phase is dried over sodium sulfate and the solvents are distilled off. The residue is purified by chromatography on silica gel (hexane/ethyl acetate), yielding 4"-desoxy-4"-epi-N-ethylamino-avermectin B1.

Example P1.3: 4"-Desoxy-4"-epi-N-(isopropoxycarbonyl-methyl)-amino-avermectin B1

300 mg of 4"-desoxy-4"-epi-amino-avermectin B1 are dissolved in 3 ml of ethyl acetate. 620 mg of isopropyl bromoacetate and 3 ml of sodium bicarbonate (1N in water) are added. The mixture is stirred vigorously at room temperature for 18 hours. The phases are then separated; the organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is purified by chromatography on silica gel (hexane/ethyl acetate), yielding 4"-desoxy-4"-epi-N-(isopropoxy-carbonyl-methyl)-amino-avermectin B1.

Example P2.1: 4"-Desoxy-4"-epi-(N-methyl-N-1-propen-3-yl-amino)-avermectin B1

600 mg of 4"-desoxy-4"-epi-methylamino-avermectin B1 are dissolved in 6 ml of ethyl acetate. 0.56 ml of allyl bromide and 6 ml of sodium bicarbonate (1N in water) are added. The mixture is stirred vigorously at 60°C for 18 hours, then cooled. The phases are then separated; the organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is purified by chromatography on silica gel (hexane/ethyl acetate), yielding 4"-desoxy-4"-epi-(N-methyl-N-1-propen-3-yl-amino)-avermectin B1.

Example P2.2: 4"-Desoxy-4"-epi-(N-2-hydroxyethyl-N-methylamino)-avermectin B1

Step 1: 4.55 g of 4"-desoxy-4"-epi-methylamino-avermectin B1 are dissolved in 45 ml of ethyl acetate. 8.6 g of ethyl bromoacetate and 45 ml of sodium bicarbonate (1N in water) are added. The mixture is stirred vigorously at 60°C for 18 hours, then cooled. The phases are then separated; the organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is purified by chromatography on silica gel (hexane/ethyl acetate), yielding 4"-desoxy-4"-epi-(N-methyl-N-ethoxycarbonylmethyl-amino)-avermectin B1.

Step 2: 300 mg of 4"-desoxy-4"-epi-(N-methyl-N-ethoxycarbonylmethyl-amino)-avermectin B1 are dissolved in 6 ml of toluene. With stirring at room temperature, 1.3 ml of diisobutylaluminium hydride (1.2 mol/l in toluene) are added. After 15 minutes, extraction is carried out with ethyl acetate and saturated ammonium chloride solution. The phases are then separated; the organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is purified by chromatography on silica gel (ethyl acetate/methanol), yielding 4"-desoxy-4"-epi-(N-2-hydroxyethyl-N-methylamino)-avermectin B1.

Example P2.3: 4"-Desoxy-4"-epi-(N-isopropyl-N-methylamino)-avermectin B1

2.0 g of 4"-desoxy-4"-epi-isopropylamino-avermectin B1 are dissolved in 20 ml of ethyl acetate. 4 ml of methyl iodide and 20 ml of sodium bicarbonate (1N in water) are added and the mixture is stirred vigorously at 60°C for 14 hours, then cooled. The phases are then separated; the organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is purified by chromatography on silica gel (ethyl acetate), yielding 4"-desoxy-4"-epi-(N-isopropyl-N-methylamino)-avermectin B1.

Example P2.4: 4"-Desoxy-4"-epi-(N-isopropyl-N-methylamino)-avermectin B1

9.14 g of 4"-desoxy-4"-epi-isopropylamino-avermectin B1 are dissolved in 100 ml of methanol. 15 ml of pivalic acid and 25 ml of formaldehyde solution (37% in water) are added. 0.7 g of sodium cyanoborohydride is then added. The mixture is stirred at room temperature for 1 hour, then the methanol is evaporated off *in vacuo* and the residue is extracted with ethyl acetate and saturated sodium bicarbonate solution. The phases are then separated; the organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is purified by chromatography on silica gel (ethyl acetate/methanol), yielding 4"-desoxy-4"-epi-(N-isopropyl-N-methylamino)-avermectin B1.

Example P2.5: 4"-Desoxy-4"-epi-(N-carboxymethyl-N-methylamino)-avermectin B1

Step 1: 10 g of 4"-desoxy-4"-epi-methylamino-avermectin B1 are dissolved in 100 ml of ethyl acetate. 15.6 g of benzyl bromoacetate and 100 ml of sodium bicarbonate (1N in water) are added. The mixture is stirred vigorously at 60°C for 5 days, then cooled. The phases are then separated; the organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is purified by chromatography on silica gel (hexane/ethyl acetate), yielding 4"-desoxy-4"-epi-(N-benzyloxycarbonyl-methyl-N-methylamino)-avermectin B1.

Step 2: 7.8 g of 4"-desoxy-4"-epi-(N-benzyloxycarbonyl-methyl-N-methylamino)-avermectin B1 are dissolved in 100 ml of tetrahydrofuran. 780 mg of palladium (5% on carbon) are added and hydrogenation is carried out at normal pressure and room temperature. After one hour the absorption of hydrogen has ceased. The mixture is filtered over Celite and the solvent is evaporated off, yielding 4"-desoxy-4"-epi-(N-carboxymethyl-N-methylamino)-avermectin B1.

Example P2.6: 4"-Desoxy-4"-epi-(N-carboxymethyl-N-methylamino)-avermectin B1

Step 1: 15 g of 4"-desoxy-4"-epi-methylamino-avermectin B1 are dissolved in 120 ml of ethyl acetate. 26 g of methyl bromoacetate and 120 ml of sodium bicarbonate (1N in water) are added. The mixture is stirred vigorously at 60°C for 5 days, then cooled. The phases are then separated; the organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is purified by chromatography on silica gel (hexane/ethyl acetate), yielding 4"-desoxy-4"-epi-(N-methoxycarbonyl-methyl-amino-N-methyl)-avermectin B1.

Step 2: 10 g of 4"-desoxy-4"-epi-(N-methoxycarbonylmethyl-amino-N-methyl)-avermectin B1 are dissolved in 90 ml of tetrahydrofuran. 10 ml of water and 440 mg of lithium hydroxide monohydrate are added and stirring is carried out at room temperature for 14 hours. Extraction is then carried out with water and diethyl ether, and the aqueous phase is separated off and lyophilised. The residue is extracted with ethyl acetate and citric acid (10% in water); the organic phase is dried over sodium sulfate and the solvent is distilled off, yielding 4"-desoxy-4"-epi-(N-carboxymethyl-N-methylamino)-avermectin B1.

Example P2.7: 4"-Desoxy-4"-epi-(N-ethyl-N-methylamino)-avermectin B1

8.0 g of 4"-desoxy-4"-epi-N-methylamino-avermectin B1 are dissolved in 50 ml of ethyl acetate. 15 ml of ethyl iodide and 50 ml of sodium bicarbonate (1N in water) are added. The mixture is stirred vigorously at 60°C for 2 days. The phases are then separated; the organic phase is dried over sodium sulfate and the solvents are distilled off. The residue is purified

by chromatography on silica gel (ethyl acetate/methanol), yielding 4"-desoxy-4"-epi-(N-ethyl-N-methylamino)-avermectin B1.

Example P2.8: 4"-Desoxy-4"-epi-[N-[(1-benzyloxycarbonyl-ethylcarbamoyl)-methyl]-N-methylamino]-avermectin B1

500 mg of 4"-desoxy-4"-epi-(N-carboxymethyl-N-methylamino)-avermectin B are dissolved in 5 ml of ethyl acetate, then 170 mg of L-alanine benzyl ester, 72 mg of 1-hydroxy-7-aza-benzotriazole and 110 mg of N,N-dicyclohexylcarbodiimide are added. Stirring is carried out at room temperature for 7 days. The mixture is then extracted with ethyl acetate and sodium bicarbonate (1N in water); the organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is purified by chromatography on silica gel (hexane/ethyl acetate), yielding 4"-desoxy-4"-epi-[N-[(1-benzyloxycarbonyl-ethylcarbamoyl)-methyl]-N-methylamino]-avermectin B1.

Example P2.9: 4"-Desoxy-4"-epi-[N-[(1-carboxy-ethylcarbamoyl)-methyl]-N-methylamino]-avermectin B1

160 mg of 4"-desoxy-4"-epi-[N-[(1-benzyloxycarbonyl-ethylcarbamoyl)-methyl]-N-methylamino]-avermectin B1 are dissolved in 10 ml of tetrahydrofuran. 50 mg of palladium (5% on carbon) are added and hydrogenation is carried out at normal pressure and room temperature. After 3 hours the absorption of hydrogen has ceased. The mixture is filtered over Celite and the solvent is evaporated off, yielding 4"-desoxy-4"-epi-[N-[(1-carboxy-ethylcarbamoyl)-methyl]-N-methylamino]-avermectin B1.

Example P2.10: 4"-Desoxy-4"-epi-[N-(2-hydroxy-2-methyl-propyl)-N-methylamino]-avermectin B1

300 mg of 4"-desoxy-4"-epi-(N-methyl-N-ethoxycarbonylmethyl-amino)-avermectin B1 (Step 1 from P2.2) are dissolved in 6 ml of tetrahydrofuran. With stirring at room temperature, 0.64 ml of methylmagnesium bromide (3 mol/l in diethyl ether) is added. After one hour, extraction is carried out with ethyl acetate and saturated ammonium chloride solution. The phases are then separated; the organic phase is dried over sodium sulfate and the solvents are distilled off. The residue is purified by chromatography on silica gel (ethyl acetate/methanol), yielding 4"-desoxy-4"-epi-[N-(2-hydroxy-2-methyl-propyl)-N-methylamino]-avermectin B1.

Example P3.1: 4"-Desoxy-4"-epi-[N,N-bis(1-phenyl-1-propen-3-yl)amino]-avermectin B1

3.48 g of 4"-desoxy-4"-epi-amino-avermectin B1 are dissolved in 40 ml of ethyl acetate. 4.62 g of 3-bromo-1-phenyl-1-propene and 40 ml of sodium bicarbonate (1N in water) are added. The mixture is stirred vigorously at 60°C for 3 days and then cooled. The phases are then separated; the organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is purified by chromatography on silica gel (hexane/ethyl acetate) yielding 4"-desoxy-4"-epi-[N,N-bis(1-phenyl-1-propen-3-yl)amino]-avermectin B1.

Example P3.2: 4"-Desoxy-4"-epi-(azetidin-1-yl)-avermectin B1

300 mg of 4"-desoxy-4"-epi-amino-avermectin B1 are dissolved in 1 ml of toluene. 0.106 ml of 1,3-dibromopropane and 1 ml of sodium bicarbonate (1N in water) are added. The mixture is stirred vigorously at 90°C for 24 hours, then cooled. The phases are then separated; the organic phase is dried over sodium sulfate and the solvent is distilled off. The residue is purified by chromatography on silica gel (ethyl acetate/methanol), yielding 4"-desoxy-4"-epi-(azetidin-1-yl)-avermectin B1.

Example P3.3: 4"-Desoxy-4"-epi-[N,N-bis(3,3-dimethyl-butyl)amino]-avermectin B1

0.87 g of 4"-desoxy-4"-epi-amino-avermectin B1 is dissolved in 10 ml of tetrahydrofuran. 1 ml of pivalic acid, 0.1 ml of water and 0.60 g of 3,3-dimethylbutyraldehyde are added. 0.38 g of sodium cyanoborohydride is then added. The mixture is stirred at room temperature for 14 hours. Extraction is then carried out with ethyl acetate and sodium bicarbonate (1N in water); the organic phase is dried over sodium sulfate and the solvents are distilled off. The residue is purified by chromatography on silica gel (hexane/ethyl acetate), yielding 4"-desoxy-4"-epi-[N,N-bis(3,3-dimethyl-butyl)amino]-avermectin B1.

Preparation of salts of formula (I)

Example P4.1: Preparation of 4"-deoxy-4"-epi-N,N-dimethylammonium-avermectin B1 benzoate

500 mg of 4"-deoxy-4"-epi-N,N-dimethylamino-avermectin B1 and 67 mg of benzoic acid are dissolved in 5 ml of dichloromethane. The solvent is evaporated off and the residue is suspended in diethyl ether and filtered off over a glass frit. The filtration residue is washed with diethyl ether and dried on the frit in a stream of air. 4"-Deoxy-4"-epi-N,N-dimethylammonium-avermectin B1 benzoate is obtained.

Example P4.2: Preparation of 4"-deoxy-4"-epi-N,N-dimethylammonium-avermectin B1 maleate (1 : 1)

500 mg of 4"-deoxy-4"-epi-N,N-dimethylamino-avermectin B1 and 64 mg of maleic acid are dissolved in 5 ml of dichloromethane. The solvent is evaporated off and the residue is suspended in diethyl ether and filtered off over a glass frit. The filtration residue is washed with diethyl ether and dried on the frit in a stream of air. 4"-Deoxy-4"-epi-N,N-dimethylammonium-avermectin B1 maleate is obtained.

Example P4.3: Preparation of 4"-deoxy-4"-epi-N,N-dimethylammonium-avermectin B1 salicylate

500 mg of 4"-deoxy-4"-epi-N,N-dimethylamino-avermectin B1 and 76 mg of salicylic acid are dissolved in 5 ml of dichloromethane. The solvent is evaporated off and the residue is suspended in diethyl ether and filtered off over a glass frit. The filtration residue is washed with diethyl ether and dried on the frit in a stream of air. 4"-Deoxy-4"-epi-N,N-dimethylammonium-avermectin B1 salicylate is obtained.

Example P4.4: Preparation of 4"-deoxy-4"-epi-N,N-dimethylammonium-avermectin B1 citrate (1 : 1)

500 mg of 4"-deoxy-4"-epi-N,N-dimethylamino-avermectin B1 and 106 mg of citric acid are dissolved in 5 ml of dichloromethane. The solvent is evaporated off and the residue is suspended in diethyl ether and filtered off over a glass frit. The filtration residue is washed with diethyl ether and dried on the frit in a stream of air. 4"-Deoxy-4"-epi-N,N-dimethylammonium-avermectin B1 citrate is obtained.

Example P4.5: Preparation of 4"-deoxy-4"-epi-N,N-dimethylammonium-avermectin B1 benzenesulfonate

500 mg of 4"-deoxy-4"-epi-N,N-dimethylamino-avermectin B1 and 87 mg of benzenesulfonic acid are dissolved in 5 ml of acetonitrile. The solvent is evaporated off and the residue is suspended in diethyl ether and filtered off over a glass frit. The filtration residue is washed with diethyl ether and dried on the frit in a stream of air. 4"-Deoxy-4"-epi-N,N-dimethylammonium-avermectin B1 benzenesulfonate is obtained.

Example P5.1: Preparation of 4"-deoxy-4"-epi-N-isopropylammonium-avermectin B1 benzoate

300 mg of 4"-deoxy-4"-epi-N-isopropylamino-avermectin B1 and 38 mg of benzoic acid are dissolved in 1 ml of acetonitrile. The solvent is evaporated off and the residue is suspended in a small amount of hexane and filtered off over a glass frit. The filtration

residue is dried on the frit in a stream of air. 4"-Deoxy-4"-epi-N-isopropylammonium-avermectin B1 benzoate is obtained.

Example P5.2: Preparation of 4"-deoxy-4"-epi-N-isopropylammonium-avermectin B1 maleate (1 : 1)

300 mg of 4"-deoxy-4"-epi-N-isopropylamino-avermectin B1 and 36 mg of maleic acid are dissolved in 1 ml of acetonitrile. 3 ml of toluene are added and then 20 ml of hexane. The mixture is filtered over a glass frit, washed with hexane and dried on the frit in a stream of air. 4"-Deoxy-4"-epi-N-isopropylammonium-avermectin B1 maleate is obtained.

Example P5.3: Preparation of 4"-deoxy-4"-epi-N-isopropylammonium-avermectin B1 salicylate

300 mg of 4"-deoxy-4"-epi-N-isopropylamino-avermectin B1 and 43 mg of salicylic acid are dissolved in 1 ml of acetonitrile. 3 ml of toluene are added and then 20 ml of hexane. The mixture is filtered over a glass frit, washed with hexane and dried on the frit in a stream of air. 4"-Deoxy-4"-epi-N-isopropylammonium-avermectin B1 salicylate is obtained.

Example P5.4: Preparation of 4"-deoxy-4"-epi-N-isopropylammonium-avermectin B1 citrate (1 : 1)

300 mg of 4"-deoxy-4"-epi-N-isopropylamino-avermectin B1 and 65 mg of citric acid are dissolved in 1 ml of acetonitrile. 3 ml of toluene are added and then 20 ml of hexane. The mixture is filtered over a glass frit, washed with hexane and dried on the frit in a stream of air. 4"-Deoxy-4"-epi-N-isopropylammonium-avermectin B1 citrate is obtained.

Example P5.5: Preparation of 4"-deoxy-4"-epi-N-isopropylammonium-avermectin B1 benzenesulfonate










300 mg of 4"-deoxy-4"-epi-N-isopropylamino-avermectin B1 and 49 mg of benzenesulfonic acid are dissolved in 1 ml of acetonitrile. 3 ml of toluene are added and then 20 ml of hexane. The mixture is filtered over a glass frit, washed with hexane and dried on the frit in a stream of air. 4"-Deoxy-4"-epi-N-isopropylammonium-avermectin B1 citrate is obtained.

Example P6.1: Analogously to the above Preparation Examples it is also possible to prepare the compounds listed in Tables 1 to 60.








Table A: Compounds of formula (I) can be formed especially with the following acids:

No.	
-----	--











No.	
A.1)	 benzoic acid
A.2)	cis-HOOC-CH=CHCOOH maleic acid
A.3)	trans-HOOC-CH=CHCOOH fumaric acid
A.4)	 2-hydroxybenzoic acid, salicylic acid
A.5)	HOOC-CH(OH)CH <sub>2</sub> COOH malic acid
A.6)	 benzenesulfonic acid
A.7)	 barbituric acid
A.8)	 2-ethylbutyric acid
A.9)	HOOC-CH(SH)CH <sub>2</sub> COOH thiomalic acid
A.10)	 3,5-dihydroxy-benzoic acid
A.11)	 trimesic acid
A.12)	 D-(-)-quinic acid
A.13)	 2-bromo-benzoic acid










No.	
A.14)	<div data-bbox="508 373 678 499"></div> <div data-bbox="841 394 1138 432">2-phenyl-benzoic acid</div>
A.15)	<div data-bbox="492 583 708 636"></div> <div data-bbox="841 569 1162 606">3,3'-thiodipropionic acid</div>
A.16)	<div data-bbox="524 716 662 842"></div> <div data-bbox="841 743 1243 781">naphthalene-1-carboxylic acid</div>
A.17)	<div data-bbox="521 873 695 1020"></div> <div data-bbox="841 915 1102 953">5-sulfosalicylic acid</div>
A.18)	<div data-bbox="508 1056 691 1182"></div> <div data-bbox="841 1077 1227 1115">2-methoxy-phenylacetic acid</div>
A.19)	<div data-bbox="500 1224 699 1350"></div> <div data-bbox="841 1251 1268 1289">benzene-1,2,4-tricarboxylic acid</div>
A.20)	<div data-bbox="540 1392 659 1518"></div> <div data-bbox="841 1419 1154 1457">3-hydroxy-benzoic acid</div>
A.21)	<div data-bbox="492 1591 708 1675"></div> <div data-bbox="841 1591 1049 1629">D-gluconic acid</div>
A.22)	<div data-bbox="521 1734 667 1871"></div> <div data-bbox="841 1766 1179 1803">4,5-dichloro-phthalic acid</div>










No.	
A.23)	$\text{HOOC}-(\text{CH}_2)_5\text{CH}_3$ n-hexanoic acid (caproic acid)
A.24)	$\text{HOOC}-(\text{CH}_2)_7\text{CH}_3$ n-heptanoic acid (oenanthic acid)
A.25)	$\text{HOOC}-(\text{CH}_2)_8\text{CH}_3$ n-octanoic acid (caprylic acid)
A.26)	$\text{HOOC}-(\text{CH}_2)_{16}\text{CH}_3$ stearic acid
A.27)	$\text{HOOC}-(\text{CH}_2)_{14}\text{CH}_3$ palmitic acid
A.28)	 2,2'-dihydroxy-1,1'-dinaphthylmethane-3,3'-4,4'-methylene-bis(3-hydroxy-2-naphthoic acid) embonic acid
A.29)	 4-methoxy-phenylacetic acid homoanistic acid
A.30)	 2-anisic acid (2-methoxy-benzoic acid)
A.31)	 adamantane-1-carboxylic acid
A.32)	 pyridine-3,4-dicarboxylic acid
A.33)	 3,4-dihydroxy-benzoic acid
A.34)	 1-hydroxy-2-naphthoic acid (1-naphthol-2-carboxylic acid)

No.	
A.35)	<div data-bbox="500 405 711 468"></div> <div data-bbox="846 401 1117 478">2,2'-oxydiacetic acid (diglycolic acid)</div>
A.36)	<div data-bbox="500 573 690 646"></div> <div data-bbox="846 569 1117 600">O-ethyl-glycolic acid</div>
A.37)	<div data-bbox="500 741 711 825"></div> <div data-bbox="846 741 1279 819">(2-naphthylthio)-acetic acid (S-(2-naphthyl)-thioglycolic acid)</div>
A.38)	<div data-bbox="500 909 711 993"></div> <div data-bbox="846 909 1166 940">2-naphthoxy-acetic acid</div>
A.39)	<div data-bbox="492 1098 719 1150"></div> <div data-bbox="846 1083 1149 1115">perfluoro-octanoic acid</div>
A.40)	<div data-bbox="508 1224 695 1350"></div> <div data-bbox="846 1251 1008 1283">p-toluic acid</div>
A.41)	<div data-bbox="500 1413 711 1518"></div> <div data-bbox="846 1419 1198 1451">cyclohexanepropionic acid</div>
A.42)	<div data-bbox="516 1560 695 1707"></div> <div data-bbox="846 1587 1365 1675">2,6-dihydroxypyridine-4-carboxylic acid (citrazinic acid)</div>
A.43)	<div data-bbox="500 1770 690 1843"></div> <div data-bbox="846 1766 1170 1797">3-methoxypropionic acid</div>









No.	
A.44)	<div data-bbox="493 359 704 537"></div> <div data-bbox="841 401 1235 485">3,4,5-trihydroxy-benzoic acid (gallic acid)</div>
A.45)	<div data-bbox="521 569 672 688"></div> <div data-bbox="841 590 1170 674">pyromucic acid (furan-2-carboxylic acid)</div>
A.46)	<div data-bbox="521 726 688 873"></div> <div data-bbox="841 758 1138 842">2-methyl-benzoic acid (o-toluic acid)</div>
A.47)	<div data-bbox="488 947 721 989"></div> <div data-bbox="841 926 1268 957">3,6,9-trioxa-undecanedioic acid</div>
A.48)	<div data-bbox="496 1094 704 1178"></div> <div data-bbox="841 1094 1317 1178">3-(4-methoxyphenyl)-propionic acid (p-methoxy-hydrocinnamic acid)</div>
A.49)	<div data-bbox="841 1262 1357 1304">3-(3,4-dihydroxyphenyl)-propionic acid</div>
A.50)	<div data-bbox="542 1314 667 1457"></div> <div data-bbox="841 1346 1138 1430">O-acetyl-salicylic acid (aspirin)</div>
A.51)	<div data-bbox="542 1486 656 1625"></div> <div data-bbox="841 1514 1127 1545">3-fluoro-benzoic acid</div>
A.52)	<div data-bbox="521 1667 678 1793"></div> <div data-bbox="841 1682 1208 1724">cyclohexanecarboxylic acid</div>










No.	
A.53)	<div data-bbox="500 373 698 504"></div> <div data-bbox="842 396 1281 480">5-chloro-2-hydroxy-benzoic acid (5-chloro-salicylic acid)</div>
A.54)	<div data-bbox="503 541 698 672"></div> <div data-bbox="842 569 1192 653">2,5-dimethyl-benzoic acid (p-xylic acid)</div>
A.55)	<div data-bbox="516 709 685 850"></div> <div data-bbox="842 741 1245 825">3,4,5-trimethoxy-benzoic acid (trimethylgallic acid)</div>
A.56)	<div data-bbox="487 877 714 997"></div> <div data-bbox="842 911 1221 945">2,4,6-trimethyl-benzoic acid</div>
A.57)	<div data-bbox="435 1018 766 1134"></div> <div data-bbox="842 1052 1167 1085">3-phenoxy-benzoic acid</div>
A.58)	<div data-bbox="527 1165 673 1302"></div> <div data-bbox="842 1192 1127 1226">4-phenyl-butyric acid</div>
A.59)	<div data-bbox="495 1344 698 1459"></div> <div data-bbox="842 1362 1245 1396">3-trifluoromethyl-benzoic acid</div>
A.60)	<div data-bbox="506 1507 686 1648"></div> <div data-bbox="842 1535 1318 1568">terephthalic acid monomethyl ester</div>
A.61)	<div data-bbox="527 1680 682 1816"></div> <div data-bbox="842 1707 1224 1740">o-hydroxy-phenylacetic acid</div>

No.	
A.62)	<div data-bbox="527 367 657 514"></div> <div data-bbox="836 399 1047 430">isophthalic acid</div>
A.63)	<div data-bbox="495 546 690 682"></div> <div data-bbox="836 577 1226 609">2,4,6-trihydroxy-benzoic acid</div>
A.64)	<div data-bbox="503 724 682 850"></div> <div data-bbox="836 745 1226 777">trifluoromethanesulfonic acid</div>
A.65)	<div data-bbox="527 892 657 1018"></div> <div data-bbox="836 913 1153 997">2-methyl-propionic acid (isobutyric acid)</div>
A.66)	<div data-bbox="511 1060 665 1197"></div> <div data-bbox="836 1081 1209 1113">2,4-dimethoxy-benzoic acid</div>
A.67)	<div data-bbox="495 1249 682 1344"></div> <div data-bbox="836 1249 1169 1333">2-thienylacetic acid (thiophene-2-acetic acid)</div>
A.68)	<div data-bbox="511 1407 690 1543"></div> <div data-bbox="836 1428 1209 1512">3,4-dimethoxy-benzoic acid (veratric acid)</div>
A.69)	<div data-bbox="511 1575 682 1711"></div> <div data-bbox="836 1596 1339 1627">2,2-bis(hydroxymethyl)-propionic acid</div>
A.70)	<div data-bbox="519 1743 673 1879"></div> <div data-bbox="836 1764 1185 1795">2-fluoro-phenylacetic acid</div>










No.	
A.71)	<div data-bbox="511 367 690 493"></div> <div data-bbox="849 394 1131 426">2-methyl-butyric acid</div>
A.72)	<div data-bbox="511 546 706 661"></div> <div data-bbox="849 562 1109 594">hydroxy-acetic acid</div>
A.73)	<div data-bbox="560 703 641 850"></div> <div data-bbox="849 735 1203 766">4-chloro-phenylacetic acid</div>
A.74)	<div data-bbox="519 871 682 1018"></div> <div data-bbox="849 907 1170 984">2-mercaptobenzoic acid (thiosalicylic acid)</div>
A.75)	<div data-bbox="503 1039 706 1186"></div> <div data-bbox="849 1075 1289 1152">(±)-2-hydroxyphenyl-acetic acid (DL-mandelic acid)</div>
A.76)	<div data-bbox="511 1207 682 1354"></div> <div data-bbox="849 1245 1398 1276">2,4-dihydropyrimidine-6-carboxylic acid</div>
A.77)	<div data-bbox="495 1396 722 1522"> H<sub>2</sub>O</div> <div data-bbox="849 1415 1154 1541">toluene-4-sulfonic acid (p-toluene-sulfonic acid)</div>
A.78)	<div data-bbox="519 1564 682 1711"></div> <div data-bbox="849 1602 1196 1633">2-chloro-phenylacetic acid</div>
A.79)	<div data-bbox="544 1732 649 1879"></div> <div data-bbox="849 1772 1180 1803">2,4-dichloro-benzoic acid</div>





No.	
A.80)	2,6-dichloro-benzoic acid
A.81)	 <chem>CC(S)C(=O)O</chem> 2-mercapto-propionic acid (thiolactic acid)
A.82)	 <chem>O=C(O)c1ccccc1Cl</chem> 2-chloro-benzoic acid
A.83)	 <chem>CS(=O)(=O)O</chem> methanesulfonic acid
A.84)	 <chem>CCS(=O)(=O)O</chem> ethanesulfonic acid (ethyl-sulfuric acid)
A.85)	 <chem>OC(=O)CCCOc1ccccc1</chem> 4-phenoxy-butyric acid
A.86)	 <chem>CC(C)(C)c1ccc(cc1)C(=O)O</chem> 4-tert-butyl-benzoic acid
A.87)	 <chem>O=C(O)c1cc2c(c1)OCO2</chem> 3,4-methylenedioxy-benzoic acid
A.88)	 <chem>OC(=O)CCSSCC(=O)O</chem> bis(2-carboxyethyl) disulfide

No.	
A.89)	<div data-bbox="524 384 667 506"></div> <div data-bbox="841 394 1125 485">pivalic acid (trimethylacetic acid)</div>
A.90)	<div data-bbox="500 548 703 674"></div> <div data-bbox="841 569 1133 604">nicotinic acid N-oxide</div>
A.91)	<div data-bbox="508 730 683 835"></div> <div data-bbox="841 741 995 777">acrylic acid</div>
A.92)	<div data-bbox="532 884 678 1024"></div> <div data-bbox="841 915 1230 995">3-benzoylpropionic acid (4-oxo-4-phenyl-butyric acid)</div>
A.93)	<div data-bbox="521 1056 699 1199"></div> <div data-bbox="841 1083 1404 1119">(1R)-(-)-camphor-10-sulfonic acid hydrate</div>
A.94)	<div data-bbox="548 1224 651 1367"></div> <div data-bbox="841 1255 1247 1291">2-chloro-4-fluoro-benzoic acid</div>
A.95)	<div data-bbox="524 1392 678 1535"></div> <div data-bbox="841 1423 1214 1459">3,5-dimethoxy-benzoic acid</div>
A.96)	<div data-bbox="513 1560 695 1703"></div> <div data-bbox="841 1598 1105 1633">2-sulfobenzoic acid</div>
A.97)	<div data-bbox="513 1745 686 1871"></div> <div data-bbox="841 1766 1052 1801">sulfoacetic acid</div>

No.	
A.98)	<div data-bbox="521 373 678 499"></div> <div data-bbox="846 401 1252 432">2-chloro-6-fluoro-benzoic acid</div>
A.99)	<div data-bbox="548 541 651 678"></div> <div data-bbox="846 573 1211 604">2,4-dihydroxy-benzoic acid</div>
A.100)	<div data-bbox="505 741 683 825"></div> <div data-bbox="846 747 1105 779">methoxyacetic acid</div>
A.101)	<div data-bbox="488 888 716 1014"></div> <div data-bbox="846 915 1333 947">2,4,6-trimethyl-benzenesulfonic acid</div>
A.102)	<div data-bbox="500 1066 708 1171"></div> <div data-bbox="846 1083 1000 1115">tartaric acid</div>
A.103)	<div data-bbox="500 1234 691 1350"></div> <div data-bbox="846 1251 1203 1283">xanthene-9-carboxylic acid</div>
A.104)	<div data-bbox="500 1423 691 1497"></div> <div data-bbox="846 1423 1065 1507">4-pentenoic acid (allylacetic acid)</div>
A.105)	<div data-bbox="508 1560 678 1728"></div> <div data-bbox="846 1598 1097 1629">5-sulfosalicylic acid</div>
A.106)	<div data-bbox="500 1791 683 1864"></div> <div data-bbox="846 1791 1040 1822">vinylacetic acid</div>

No.	
A.107)	 <div>2-butynedioic acid (acetylenedicarboxylic acid)</div>
A.108)	 <div>2-oxo-propionic acid (pyruvic acid)</div>
A.109)	 <div>cyclohexylacetic acid</div>
A.110)	 <div>2-hydroxyisobutyric acid</div>
A.111)	 <div>nicotinic acid</div>
A.112)	 <div>6-chloro-nicotinic acid</div>
A.113)	 <div>isonicotinic acid</div>
A.114)	 <div>picolinic acid</div>
A.115)	 <div>pyrazinecarboxylic acid</div>
A.116)	<div>HOOC-COOH</div> <div>oxalic acid</div>
A.117)	<div>CH<sub>3</sub>CH<sub>2</sub>COOH</div> <div>propionic acid</div>
A.118)	<div>CF<sub>3</sub>CF<sub>2</sub>COOH</div> <div>pentafluoropropionic acid</div>
A.119)	<div>CH<sub>3</sub>(CH<sub>2</sub>)<sub>2</sub>COOH</div> <div>butyric acid</div>
A.120)	<div>CF<sub>3</sub>CF<sub>2</sub>CF<sub>2</sub>COOH</div> <div>heptafluorobutyric acid</div>
A.121)	<div>CH<sub>3</sub>(CH<sub>2</sub>)<sub>3</sub>COOH</div> <div>valeric acid</div>

No.	
A.122)	<div style="display: flex; align-items: center;"> <div style="text-align: center; margin-right: 20px;"> <math display="block">  \begin{array}{c}  \text{COOH} \\    \\  \text{HOOC}-\text{CH}-\text{CH}_2-\text{COOH} \\    \\  \text{OH}  \end{array}  </math> </div> <div>citric acid</div> </div>
A.123)	HOCH <sub>2</sub> CH(OH)COOH      glyceric acid
A.124)	CH <sub>3</sub> COOH      acetic acid
A.125)	ClCH <sub>2</sub> COOH      chloroacetic acid
A.126)	Cl <sub>2</sub> CHCOOH      dichloroacetic acid
A.127)	CF <sub>3</sub> COOH      trifluoroacetic acid
A.128)	FCH <sub>2</sub> COOH      fluoroacetic acid
A.129)	CH <sub>3</sub> CH(OH)COOH      lactic acid
A.130)	HOOCCH <sub>2</sub> COOH      malonic acid
A.131)	HOOC-(CH <sub>2</sub> ) <sub>2</sub> COOH      succinic acid
A.132)	HOOC-(CH <sub>2</sub> ) <sub>3</sub> COOH      glutaric acid
A.133)	HOOC-(CH <sub>2</sub> ) <sub>4</sub> COOH      adipic acid
A.134)	HOOC-(CH <sub>2</sub> ) <sub>5</sub> COOH      pimelic acid
A.135)	HOOC-(CH <sub>2</sub> ) <sub>6</sub> COOH      suberic acid
A.136)	HOOC-(CH <sub>2</sub> ) <sub>7</sub> COOH      azelaic acid
A.137)	HOOC-(CH <sub>2</sub> ) <sub>8</sub> COOH      sebacic acid
A.138)	<div style="display: flex; align-items: center;"> <div style="text-align: center; margin-right: 20px;">  </div> <div>phthalic acid</div> </div>
A.139)	<div style="display: flex; align-items: center;"> <div style="text-align: center; margin-right: 20px;">  </div> <div>terephthalic acid</div> </div>
A.140)	H <sub>3</sub> PO <sub>4</sub> phosphoric acid
A.141)	H <sub>2</sub> SO <sub>4</sub> sulfuric acid
A.142)	HCl      hydrochloric acid
A.143)	HBr      hydrobromic acid
A.144)	HI      hydriodic acid
A.145)	HNO <sub>3</sub> nitric acid
A.146)	HClO <sub>4</sub> perchloric acid
A.147)	CH <sub>3</sub> C(=O)-CH <sub>2</sub> -COOH      acetoacetic acid
A.148)	NC-CH <sub>2</sub> -COOH      cyanoacetic acid



No.	
A.149)	<div style="display: flex; align-items: center; justify-content: center;">  <div style="margin-left: 20px;">tetrahydrofuran-2-carboxylic acid</div> </div>
A.150)	<div style="display: flex; align-items: center; justify-content: center;"> <math>\text{CH}\equiv\text{C}-\text{COOH}</math> <div style="margin-left: 20px;">propionic acid</div> </div>
A.151)	<div style="display: flex; align-items: center; justify-content: center;"> <math>\text{H}_2\text{C}=\text{C}(\text{CH}_3)-\text{COOH}</math> <div style="margin-left: 20px;">methacrylic acid</div> </div>
A.152)	<div style="display: flex; align-items: center; justify-content: center;"> <math>\text{CH}_3-\text{CH}=\text{CH}-\text{COOH}</math> <div style="margin-left: 20px;">crotonic acid</div> </div>
A.153)	<div style="display: flex; align-items: center; justify-content: center;">  <div style="margin-left: 20px;">picric acid</div> </div>

Table 1: A compound of formula (I) wherein  $R_1$  is isopropyl and  $R_2$  and  $R_3$  are hydrogen, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 2: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is methyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 3: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is ethyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 4: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is n-propyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 5: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is isopropyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 6: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is n-butyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 7: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is isobutyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 8: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is sec-butyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 9: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is tert-butyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 10: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  and  $R_3$  are ethyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 11: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is  $-\text{CH}_2\text{-CH=CH}_2$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 12: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is  $-\text{CH}_2\text{-C(=O)OC}_2\text{H}_5$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 13: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is benzyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 14: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  and  $R_3$  together are  $-\text{CH}_2\text{-CH}_2\text{-CH}_2^-$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 15: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  and  $R_3$  together are  $-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2^-$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 16: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is  $-\text{CH}_2\text{-CH}_2\text{-OH}$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

**Table 17:** A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is  $-\text{CH}_2-\text{C}(\text{CH}_3)\text{OH}$ , and  $\text{X}^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

**Table 18:** A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is hydrogen and  $R_3$  is ethyl, and  $\text{X}^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

**Table 19:** A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is hydrogen and  $R_3$  is  $-\text{CH}_2-\text{CH}=\text{CH}_2$ , and  $\text{X}^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

**Table 20:** A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is hydrogen and  $R_3$  is  $-\text{CH}_2-\text{CH}\equiv\text{CH}$ , and  $\text{X}^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

**Table 21:** A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is  $-\text{CH}_2-\text{CH}\equiv\text{CH}$ , and  $\text{X}^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.


**Table 22:** A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is hydrogen and  $R_3$  is isopropyl, and  $\text{X}^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

**Table 23:** A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is hydrogen and  $R_3$  is n-propyl, and  $\text{X}^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

**Table 24:** A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is hydrogen and  $R_3$  is n-butyl, and  $\text{X}^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

**Table 25:** A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is hydrogen and  $R_3$  is  $-\text{CH}_2-\text{CH}(\text{CH}_3)_2$ , and  $\text{X}^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.


**Table 26:** A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is hydrogen and  $R_3$  is



and  $\text{X}^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.



Table 27: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is hydrogen and  $R_3$  is



, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 28: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is hydrogen and  $R_3$  is  $-\text{CH}_2-\text{C}(=\text{O})\text{OC}_2\text{H}_5$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 29: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is hydrogen and  $R_3$  is  $-\text{CH}_2-\text{C}_6\text{H}_4-\text{O}-\text{CF}_2\text{H}$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 30: A compound of formula (I) wherein  $R_1$  is isopropyl,  $R_2$  is methyl and  $R_3$  is  $-\text{CH}_2-\text{CH}=\text{CH}-\text{C}(=\text{O})\text{OCH}_2\text{C}(=\text{O})\text{C}_6\text{H}_5$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 31: A compound of formula (I) wherein  $R_1$  is sec-butyl and  $R_2$  and  $R_3$  are hydrogen, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 32: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is methyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 33: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is ethyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 34: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is n-propyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 35: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is isopropyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 36: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is n-butyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 37: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is isobutyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 38: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is sec-butyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 39: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is tert-butyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 40: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  and  $R_3$  are ethyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 41: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is  $-\text{CH}_2\text{-CH=CH}_2$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 42: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is  $-\text{CH}_2\text{-C(=O)OC}_2\text{H}_5$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 43: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is benzyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 44: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  and  $R_3$  together are  $-\text{CH}_2\text{-CH}_2\text{-CH}_2-$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 45: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  and  $R_3$  together are  $-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2-$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 46: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is  $-\text{CH}_2\text{-CH}_2\text{-OH}$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 47: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is  $-\text{CH}_2-\text{C}(\text{CH}_3)\text{OH}$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 48: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is hydrogen and  $R_3$  is ethyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 49: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is hydrogen and  $R_3$  is  $-\text{CH}_2-\text{CH}=\text{CH}_2$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 50: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is hydrogen and  $R_3$  is  $-\text{CH}_2-\text{CH}\equiv\text{CH}$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 51: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is methyl and  $R_3$  is  $-\text{CH}_2-\text{CH}\equiv\text{CH}$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.


Table 52: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is hydrogen and  $R_3$  is isopropyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 53: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is hydrogen and  $R_3$  is n-propyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 54: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is hydrogen and  $R_3$  is n-butyl, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.


Table 55: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is hydrogen and  $R_3$  is  $-\text{CH}_2-\text{CH}(\text{CH}_3)_2$ , and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Table 56: A compound of formula (I) wherein  $R_1$  is sec-butyl,  $R_2$  is hydrogen and  $R_3$  is



and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

**Table 57:** A compound of formula (I) wherein R<sub>1</sub> is sec-butyl, R<sub>2</sub> is hydrogen and R<sub>3</sub> is



—N, and  $X^-$  in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.


**Table 58:** A compound of formula (I) wherein R<sub>1</sub> is sec-butyl, R<sub>2</sub> is hydrogen and R<sub>3</sub> is -CH<sub>2</sub>-C(=O)OC<sub>2</sub>H<sub>5</sub>, and X<sup>-</sup> in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

**Table 59:** A compound of formula (I) wherein R<sub>1</sub> is sec-butyl, R<sub>2</sub> is hydrogen and R<sub>3</sub> is -CH<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>-O-CF<sub>2</sub>H, and X<sup>-</sup> in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

**Table 60:** A compound of formula (I) wherein R<sub>1</sub> is sec-butyl, R<sub>2</sub> is methyl and R<sub>3</sub> is -CH<sub>2</sub>-CH=CH-C(=O)OCH<sub>2</sub>C(=O)C<sub>6</sub>H<sub>5</sub>, and X<sup>-</sup> in each case corresponds to the anion of one of the acids mentioned in lines A.1 to A.153 of Table A.

Tables B and C below show experimentally determined percentage contents of C, H and N in compounds of formula (I) above. Since the compounds are mixtures of avermectin derivatives B1a and B1b wherein R<sub>1</sub> is isopropyl and sec-butyl, respectively, and the proportion thereof in the mixture is variable, the Tables do not give mathematically determined values for the contents of C, H and N.

Table B: Elemental analyses of compounds of formula




No.	X-H	found content (%)		
		C	H	N
B.1	benzoic acid	65.9	8.1	1.4
B.2	Maleic acid	61.8	8.0	1.4
B.3	2-hydroxybenzoic acid, salicylic acid	62.4	7.7	1.3
B.4	benzenesulfonic acid	61.0	7.8	1.5
B.5	2-ethylbutyric acid	65.4	8.7	1.4
B.6	thiomalic acid	60.1	7.9	1.3
B.7	3,5-dihydroxy-benzoic acid	62.3	7.9	1.3
B.8	trimesic acid	61.2	7.5	1.2
B.9	D-(-)-quinic acid	60.2	8.1	1.3
B.10	2-bromo-benzoic acid	60.4	7.4	1.2
B.11	2-phenyl-benzoic acid	68.3	8.0	1.2
B.12	3,3'-thiodipropionic acid	60.9	8.1	1.3
B.13	naphthalene-1-carboxylic acid	67.6	7.9	1.2
B.14	5-sulfosalicylic acid	59.4	7.6	1.3
B.15	2-methoxy-phenylacetic acid	65.1	8.2	1.2
B.16	benzene-1,2,4-tricarboxylic acid	61.7	7.5	1.3
B.17	3-hydroxy-benzoic acid	64.0	7.7	1.3
B.18	D-gluconic acid	56.8	7.9	1.1
B.19	4,5-dichloro-phthalic acid	60.0	7.2	1.3
B.20	n-octanoic acid (caprylic acid)	65.9	8.9	1.3
B.21	2,2'-dihydroxy-1,1'-dinaphthylmethane-3,3'-4,4'-methylene-bis(3-hydroxy-2-naphthoic acid) embonic acid	64.7	7.5	1.2
B.22	4-methoxy-phenylacetic acid (homoanisic acid)	65.3	8.0	1.2
B.23	2-anisic acid (2-methoxy-benzoic acid)	65.0	7.9	1.2
B.24	adamantane-1-carboxylic acid	67.3	8.7	1.2
B.25	pyridine-3,4-dicarboxylic acid	62.8	8.2	2.9

No.	X-H	found content (%)		
		C	H	N
B.26	3,4-dihydroxy-benzoic acid	62.0	7.5	1.2
B.27	1-hydroxy-2-naphthoic acid (1-naphthol-2-carboxylic acid)	66.5	8.1	1.2
B.28	2,2'-oxydiacetic acid (diglycolic acid)	60.3	8.2	1.4
B.29	O-ethyl-glycolic acid	63.6	8.8	1.3
B.30	(2-naphthylthio)-acetic acid (S-(2-naphthyl)-thioglycolic acid)	67.1	8.2	1.2
B.31	2-naphthoxy-acetic acid	67.6	8.3	1.1
B.32	p-toluic acid	68.1	8.9	1.3
B.33	cyclohexanepropionic acid	67.4	9.1	1.3
B.34	3-methoxypropionic acid	64.0	8.7	1.4
B.35	3,4,5-trihydroxy-benzoic acid (gallic acid)	61.7	8.0	1.3
B.36	pyromucic acid (furan-2-carboxylic acid)	64.1	8.6	1.2
B.37	2-methyl-benzoic acid (o-toluic acid)	67.3	8.4	1.3
B.38	3,6,9-trioxa-undecanedioic acid	60.2	8.2	1.1
B.39	3-(4-methoxyphenyl)-propionic acid (p-methoxy-hydrocinnamic acid)	66.5	8.4	1.2
B.40	O-acetyl-salicylic acid (aspirin)	65.2	8.4	1.3
B.41	3-fluoro-benzoic acid	64.3	8.0	1.3
B.42	cyclohexanecarboxylic acid	66.6	9.0	1.4
B.43	5-chloro-2-hydroxy-benzoic acid (5-chloro-salicylic acid)	63.7	8.1	1.2
B.44	2,5-dimethyl-benzoic acid (p-xylic acid)	67.7	8.7	1.2
B.45	3,4,5-trimethoxy-benzoic acid (trimethylgallic acid)	65.9	8.6	1.2
B.46	4-phenyl-butyric acid	67.6	8.6	1.2
B.47	3-trifluoromethyl-benzoic acid	64.3	8.0	1.2
B.48	o-hydroxy-phenylacetic acid	65.0	8.3	1.2

No.	X-H	found content (%)		
		C	H	N
B.49	isophthalic acid	63.7	7.9	1.1
B.50	2,4,6-trihydroxy-benzoic acid	63.2	8.2	1.1
B.51	trifluoromethanesulfonic acid	66.6	8.9	1.4
B.52	2-methyl-propionic acid (isobutyric acid)	66.0	9.0	1.3
B.53	2-thienylacetic acid (thiophene-2-acetic acid)	65.2	8.4	1.2
B.54	3,4-dimethoxy-benzoic acid (veratric acid)	65.1	8.3	1.3
B.55	2,2-bis(hydroxymethyl)-propionic acid	64.9	8.9	1.2
B.56	2-fluoro-phenylacetic acid	66.4	8.4	1.3
B.57	2-methyl-butyric acid	66.2	9.0	1.4
B.58	hydroxy-acetic acid	61.8	8.6	1.3
B.59	4-chloro-phenylacetic acid	65.3	8.3	1.1
B.60	2-mercaptobenzoic acid (thiosalicylic acid)	63.3	8.1	1.2
B.61	(+/-)-2-hydroxyphenyl-acetic acid (DL-mandelic acid)	63.1	8.0	1.1
B.62	2,4-dihydropyrimidine-6-carboxylic acid	55.4	7.9	3.1
B.63	toluene-4-sulfonic acid (p-toluene-sulfonic acid)	61.9	8.3	1.2
B.64	2-chloro-phenylacetic acid	66.1	8.6	1.1
B.65	2,4-dichloro-benzoic acid	61.1	7.6	1.2
B.66	2-mercapto-propionic acid (thiolactic acid)	62.9	8.7	1.2
B.67	2-chloro-benzoic acid	63.3	7.8	1.1
B.68	methanesulfonic acid	59.7	8.6	1.1
B.69	ethanesulfonic acid (ethyl-sulfuric acid)	59.2	8.5	1.3
B.70	4-phenoxy-butyric acid	68.1	9.1	1.1
B.71	4-tert-butyl-benzoic acid	68.7	9.0	1.2
B.72	bis(2-carboxyethyl) disulfide	59.1	8.0	1.1
B.73	pivalic acid (trimethylacetic acid)	65.8	8.9	1.4
B.74	acrylic acid	65.4	8.8	1.3

No.	X-H	found content (%)		
		C	H	N
B.75	3-benzoylpropionic acid (4-oxo-4-phenyl-butyric acid)	67.7	8.8	1.1
B.76	(1R)-(-)-camphor-10-sulfonic acid hydrate	63.4	8.7	1.1
B.77	2-chloro-4-fluoro-benzoic acid	62.3	7.9	1.2
B.78	3,5-dimethoxy-benzoic acid	66.1	8.5	1.2
B.79	2-sulfobenzoic acid	60.3	7.9	1.2
B.80	sulfoacetic acid	59.6	8.3	1.3
B.81	2-chloro-6-fluoro-benzoic acid	61.7	7.9	1.3
B.82	2,4-dihydroxy-benzoic acid	62.6	8.2	1.3
B.83	methoxyacetic acid	63.4	8.7	1.3
B.84	tartaric acid	58.4	8.1	1.2
B.85	xanthene-9-carboxylic acid	68.3	8.3	1.1
B.86	4-pentenoic acid (allylacetic acid)	66.1	8.9	1.4
B.87	vinylacetic acid	64.9	8.6	1.3
B.88	2-butyndioic acid (acetylenedicarboxylic acid)	61.8	8.3	1.4
B.89	2-oxo-propionic acid (pyruvic acid)	62.0	8.6	1.2
B.90	cyclohexylacetic acid	66.6	9.0	1.3
B.91	2-hydroxyisobutyric acid	62.4	8.8	1.3
B.92	citric acid	59.5	7.8	1.4
B.93	adipic acid	61.1	8.1	1.2
B.94	sulfuric acid	60.7	8.6	1.8
B.95	hydrochloric acid	59.7	8.4	1.4



Table C: Elemental analyses of compounds of formula

No.	X-H	found content (%)		
		C	H	N
C.1	benzoic acid	66.3	8.2	1.5
C.2	maleic acid	61.8	8.1	1.4
C.3	2-hydroxybenzoic acid, salicylic acid	63.6	7.9	1.4
C.4	benzenesulfonic acid	61.6	7.9	1.4
C.5	citric acid	62.4	7.8	1.5

Formulation Examples for use in crop protection (% = percent by weight)Example F1: Emulsifiable concentrates

	a)	b)	c)
active ingredient	25%	40%	50%
calcium dodecylbenzenesulfonate	5%	8%	6%
castor oil polyethylene glycol ether (36 mol EO)	5%	-	-
tributylphenol polyethylene glycol ether (30 mol EO)	-	12%	4%
cyclohexanone	-	15%	20%
xylene mixture	65%	25%	20%

Mixing finely ground active ingredient and additives gives an emulsifiable concentrate which yields emulsions of the desired concentration on dilution with water.

Example F2: Solutions

	a)	b)	c)	d)
active ingredient	80%	10%	5%	95%
ethylene glycol monomethyl ether	20%	-	-	-
polyethylene glycol (MW 400)	-	70%	-	-
N-methylpyrrolid-2-one	-	20%	-	-
epoxidised coconut oil	-	-	1%	5%
benzine (boiling range: 160-190°)	-	-	94%	-

Mixing finely ground active ingredient and additives gives a solution suitable for use in the form of microdrops.

Example F3: Granules

	a)	b)	c)	d)
active ingredient	5%	10%	8%	21%
kaolin	94%	-	79%	54%
highly dispersed silicic acid	1%	-	13%	7%
attapulgate	-	90%	-	18%

The active ingredient is dissolved in dichloromethane, the solution is sprayed onto the carrier mixture and the solvent is evaporated off *in vacuo*.

Example F4: Wettable powders

	a)	b)	c)
active ingredient	25%	50%	75%
sodium lignosulfonate	5%	5%	-
sodium lauryl sulfate	3%	-	5%
sodium diisobutylphthalenesulfonate	-	6%	10%
octylphenol polyethylene glycol ether (7-8 mol EO)	-	2%	-
highly dispersed silicic acid	5%	10%	10%
kaolin	62%	27%	-

Active ingredient and additives are mixed together and the mixture is ground in a suitable mill, yielding wettable powders that can be diluted with water to form suspensions of the desired concentration.

Example F5: Emulsifiable concentrate

active ingredient	10%
octylphenol polyethylene glycol ether (4-5 mol EO)	3%
calcium dodecylbenzenesulfonate	3%
castor oil polyethylene glycol ether (36 mol EO)	4%
cyclohexanone	30%
xylene mixture	50%

Mixing finely ground active ingredient and additives gives an emulsifiable concentrate which yields emulsions of the desired concentration on dilution with water.

Example F6: Extruder granules

active ingredient	10%
sodium lignosulfonate	2%
carboxymethylcellulose	1%
kaolin	87%

Active ingredient and additives are mixed together, the mixture is ground, moistened with water, extruded and granulated and the granules are dried in a stream of air.

Example F7: Coated granules

active ingredient	3%
polyethylene glycol (MW 200)	3%
kaolin	94%

Uniform application of the finely ground active ingredient to the kaolin moistened with polyethylene glycol in a mixer yields non-dusty coated granules.

Example F8: Suspension concentrate

active ingredient	40%
ethylene glycol	10%
nonylphenol polyethylene glycol ether (15 mol EO)	6%
sodium lignosulfonate	10%
carboxymethylcellulose	1%
aqueous formaldehyde solution (37%)	0.2%
aqueous silicone oil emulsion (75%)	0.8%
water	32%

Mixing finely ground active ingredient and additives gives a suspension concentrate which yields suspensions of the desired concentration on dilution with water.

Biological Examples:

Example B1: Action against *Spodoptera littoralis*

Young soybean plants are sprayed with an aqueous emulsion spray mixture comprising 12.5 ppm of test compound and, after the spray-coating has dried, the plants are populated with 10 caterpillars of *Spodoptera littoralis* in the first stage and then placed in a plastics container. 3 days later, the percentage reduction in population and the percentage reduction in feeding damage (% activity) are determined by comparing the number of dead caterpillars and the feeding damage on the treated plants with that on untreated plants.

Example B2: Action against *Spodoptera littoralis*, systemic:

Maize seedlings are placed in the test solution. 6 days later, the leaves are cut off, placed on moist filter paper in a petri dish and infested with 12 to 15 *Spodoptera littoralis* larvae in the L<sub>1</sub> stage. 4 days later, the percentage reduction in population (% activity) is determined by comparing the number of dead caterpillars on treated plants with that on untreated plants.

Example B3: Action against *Heliothis virescens*

30-35 eggs of *Heliothis virescens*, from 0 to 24 hours old, are placed on filter paper in a petri dish on a layer of artificial nutrient. 0.8 ml of the test solution is then pipetted onto the filter papers. Evaluation is made 6 days later. The percentage reduction in population (% activity) is determined by comparing the number of dead eggs and larvae on treated plants with that on untreated plants.

Example B4: Action against *Plutella xylostella* caterpillars

Young cabbage plants are sprayed with an aqueous emulsion spray mixture comprising 12.5 ppm of test compound. After the spray-coating has dried, the cabbage plants are populated with 10 caterpillars of *Plutella xylostella* in the first stage and placed in a plastics container. Evaluation is made 3 days later. The percentage reduction in population and the percentage reduction in feeding damage (% activity) are determined by comparing the number of dead caterpillars and the feeding damage on the treated plants with that on the untreated plants.

Example B5: Action against *Frankliniella occidentalis*

Pieces of bean leaves are placed on agar in petri dishes and sprayed with test solution in a spray chamber. The leaves are then infested with a mixed population of *Frankliniella occidentalis*. Evaluation is made 10 days later. The percentage reduction (% activity) is determined by comparing the population on the treated leaves with that on untreated leaves.

Example B6: Action against *Diabrotica balteata*

Maize seedlings are sprayed with an aqueous emulsion spray mixture comprising 12.5 ppm of the test compound and, after the spray-coating has dried, the maize seedlings are populated with 10 *Diabrotica balteata* larvae in the second stage and then placed in a plastics container. 6 days later, the percentage reduction in population (% activity) is determined by comparing the number of dead larvae on the treated plants with that on untreated plants.


Example B7: Action against *Tetranychus urticae*

Young bean plants are populated with a mixed population of *Tetranychus urticae* and sprayed one day later with an aqueous emulsion spray mixture comprising 12.5 ppm of test compound. The plants are incubated for 6 days at 25°C and subsequently evaluated. The percentage reduction in population (% activity) is determined by comparing the number of dead eggs, larvae and adults on the treated plants with that on untreated plants.

The compounds of the Tables exhibit a good action in the above tests B1 to B7. For example, especially compounds B.1 to B.4, B.11, B.22, B.29, B.32, B.36, B.41, B.44, B.47, B.51, B.52, B.60, B.70, B.71, B.74, B.82, B.83, B.84, B.86, B.91, B.92 and B.94 bring about a reduction in the pest population mentioned in these tests of more than 80 %.

Patent claims:

## 1. A compound of formula



wherein

$X^-$  is an anion;

$n$  is 1, 2, 3 or 4;

$R_1$  is  $C_1$ - $C_{12}$ alkyl,  $C_3$ - $C_8$ cycloalkyl; or  $C_2$ - $C_{12}$ alkenyl;

$R_2$  is hydrogen, unsubstituted or mono- to penta-substituted  $C_1$ - $C_{12}$ alkyl, unsubstituted or mono- to penta-substituted  $C_2$ - $C_{12}$ alkenyl;

$R_3$  is hydrogen, unsubstituted or mono- to penta-substituted  $C_1$ - $C_{12}$ alkyl, unsubstituted or mono- to penta-substituted  $C_3$ - $C_{12}$ cycloalkyl, unsubstituted or mono- to penta-substituted  $C_2$ - $C_{12}$ alkenyl; unsubstituted or mono- to penta-substituted  $C_2$ - $C_{12}$ alkynyl; or

$R_2$  and  $R_3$  together are a three- to seven-membered alkylene bridge, or a four- to seven-membered alkenylene bridge wherein a  $-CH_2-$  group may have been replaced by O, S or  $NR_4$ ;

and wherein the substituents of the mentioned alkyl, alkenyl, alkynyl, alkylene, alkenylene and cycloalkyl radicals are selected from the group consisting of OH, halogen, halo- $C_1$ - $C_2$ alkyl, CN,  $NO_2$ ,  $C_2$ - $C_6$ alkynyl,  $C_3$ - $C_8$ cycloalkyl, norbornylenyl,  $C_3$ - $C_8$ cycloalkenyl;  $C_3$ - $C_8$ cycloalkenyl unsubstituted or substituted by from one to three methyl groups;

C<sub>3</sub>-C<sub>8</sub>halocycloalkyl, C<sub>1</sub>-C<sub>12</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>cycloalkoxy, C<sub>1</sub>-C<sub>12</sub>haloalkoxy, C<sub>1</sub>-C<sub>12</sub>alkylthio, C<sub>3</sub>-C<sub>8</sub>cycloalkylthio, C<sub>1</sub>-C<sub>12</sub>haloalkylthio, C<sub>1</sub>-C<sub>12</sub>alkylsulfinyl, C<sub>3</sub>-C<sub>8</sub>cycloalkylsulfinyl, C<sub>1</sub>-C<sub>12</sub>haloalkylsulfinyl, C<sub>3</sub>-C<sub>8</sub>halocycloalkylsulfinyl, C<sub>1</sub>-C<sub>12</sub>alkylsulfonyl, C<sub>3</sub>-C<sub>8</sub>cycloalkylsulfonyl, C<sub>1</sub>-C<sub>12</sub>haloalkylsulfonyl, C<sub>3</sub>-C<sub>8</sub>halocycloalkylsulfonyl, C<sub>2</sub>-C<sub>8</sub>alkenyl, C<sub>2</sub>-C<sub>8</sub>alkynyl, NH(C<sub>1</sub>-C<sub>6</sub>alkyl), N(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub>, -C(=O)R<sub>5</sub>, -NHC(=O)R<sub>6</sub>, -P(=O)(OC<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub>;

aryl, heterocyclyl, aryloxy, heterocycloxy; and also aryl, heterocyclyl, aryloxy and heterocycloxy that, depending upon the possibilities of substitution at the ring, are mono- to penta-substituted by substituents selected from the group consisting of OH, halogen, CN, NO<sub>2</sub>, C<sub>1</sub>-C<sub>12</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>1</sub>-C<sub>12</sub>haloalkyl, C<sub>1</sub>-C<sub>12</sub>alkoxy, C<sub>1</sub>-C<sub>12</sub>haloalkoxy, C<sub>1</sub>-C<sub>12</sub>alkylthio, C<sub>1</sub>-C<sub>12</sub>haloalkylthio, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkyl, dimethylamino-C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>2</sub>-C<sub>8</sub>alkenyl, C<sub>2</sub>-C<sub>8</sub>alkynyl, phenoxy, phenyl-C<sub>1</sub>-C<sub>6</sub>alkyl; phenoxy unsubstituted or substituted by from one to three substituents selected independently of one another from halogen, methoxy, trifluoromethyl and trifluoromethoxy; phenyl-C<sub>1</sub>-C<sub>6</sub>alkoxy unsubstituted or substituted in the aromatic ring by from one to three substituents selected independently of one another from halogen, methoxy, trifluoromethyl and trifluoromethoxy; phenyl-C<sub>2</sub>-C<sub>6</sub>alkenyl, phenyl-C<sub>2</sub>-C<sub>6</sub>alkynyl, methylenedioxy, -C(=O)R<sub>5</sub>, -O-C(=O)R<sub>6</sub>, -NH-C(=O)R<sub>6</sub>, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>12</sub>alkyl), N(C<sub>1</sub>-C<sub>12</sub>alkyl)<sub>2</sub>, C<sub>1</sub>-C<sub>6</sub>alkylsulfinyl, C<sub>3</sub>-C<sub>8</sub>cycloalkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>haloalkylsulfinyl, C<sub>3</sub>-C<sub>8</sub>halocycloalkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>alkylsulfonyl, C<sub>3</sub>-C<sub>8</sub>cycloalkylsulfonyl, C<sub>1</sub>-C<sub>6</sub>haloalkylsulfonyl and C<sub>3</sub>-C<sub>8</sub>halocycloalkylsulfonyl;

R<sub>4</sub> is C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>8</sub>cycloalkyl, C<sub>2</sub>-C<sub>8</sub>alkenyl, C<sub>2</sub>-C<sub>8</sub>alkynyl, benzyl or -C(=O)-R<sub>5</sub>;

R<sub>5</sub> is H, OH, SH, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>12</sub>alkyl), N(C<sub>1</sub>-C<sub>12</sub>alkyl)<sub>2</sub>, C<sub>1</sub>-C<sub>12</sub>alkyl, C<sub>1</sub>-C<sub>12</sub>haloalkyl, C<sub>1</sub>-C<sub>12</sub>alkoxy, C<sub>1</sub>-C<sub>12</sub>haloalkoxy, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>12</sub>alkylthio, C<sub>2</sub>-C<sub>8</sub>alkenyloxy, C<sub>2</sub>-C<sub>8</sub>alkynyloxy; phenyl, phenoxy, benzyloxy, -NH-phenyl, -N(C<sub>1</sub>-C<sub>6</sub>alkyl)-phenyl, NH-C<sub>1</sub>-C<sub>6</sub>alkyl-C(=O)-R<sub>7</sub>, -N(C<sub>1</sub>-C<sub>6</sub>alkyl)-C<sub>1</sub>-C<sub>6</sub>alkyl-C(=O)-R<sub>7</sub>; or phenyl, phenoxy, benzyloxy, NH-phenyl or -N(C<sub>1</sub>-C<sub>6</sub>alkyl)-phenyl substituted in the aromatic ring by from one to three substituents selected independently of one another from halogen, C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>haloalkyl and C<sub>1</sub>-C<sub>6</sub>haloalkoxy;

R<sub>6</sub> is H, C<sub>1</sub>-C<sub>12</sub>alkyl, C<sub>1</sub>-C<sub>12</sub>haloalkyl, C<sub>2</sub>-C<sub>8</sub>alkenyl, C<sub>2</sub>-C<sub>8</sub>alkynyl, phenyl, benzyl, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>12</sub>alkyl), N(C<sub>1</sub>-C<sub>12</sub>alkyl)<sub>2</sub>, -NH-phenyl or -N(C<sub>1</sub>-C<sub>12</sub>alkyl)-phenyl; and

R<sub>7</sub> is H, OH, C<sub>1</sub>-C<sub>12</sub>alkyl, C<sub>1</sub>-C<sub>12</sub>alkoxy, C<sub>1</sub>-C<sub>6</sub>alkoxy-C<sub>1</sub>-C<sub>6</sub>alkoxy, C<sub>2</sub>-C<sub>8</sub>alkenyloxy, phenyl, phenoxy, benzyloxy, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>12</sub>alkyl), N(C<sub>1</sub>-C<sub>12</sub>alkyl)<sub>2</sub>, -NH-phenyl or -N(C<sub>1</sub>-C<sub>12</sub>alkyl)-phenyl;

and, where applicable, an E/Z isomer, E/Z isomeric mixture and/or tautomer thereof,  
with the proviso that R<sub>1</sub> is not sec-butyl or isopropyl when R<sub>2</sub> is H and R<sub>3</sub> is methyl.

2. A compound according to claim 1 of formula (I) wherein R<sub>1</sub> is isopropyl or sec-butyl.
3. A pesticidal composition comprising as active ingredient at least one compound according to claim 1 of formula (I), in free form or in agrochemically acceptable salt form, and at least one adjuvant.
4. A method of controlling pests, which comprises applying a composition according to claim 3 to the pests or to the locus thereof.
5. A process for the preparation of a composition according to claim 3 comprising at least one adjuvant, which comprises intimately mixing and/or grinding the active ingredient with the adjuvant(s).
6. Use of a compound according to claim 1 of formula (I), in free form or in agrochemically acceptable salt form, in the preparation of a composition as described in claim 3.
7. Use of a composition according to claim 3 in controlling pests.
8. A method according to claim 4 for the protection of plant propagation material, which comprises treating the propagation material or the planting site of the propagation material.
9. Plant propagation material treated in accordance with the method described in claim 8.



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/02044

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07H19/01 A01N43/90

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07H A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 362 863 A (CVETOVICH RAYMOND) 8 November 1994 (1994-11-08) column 5, lines 52-60; column 6, lines 66-67; column 13, line 2: 4"-epi-amino-4"-deoxyavermectin B1 benzoate	1-9
Y	EP 0 465 121 A (MERCK & CO INC) 8 January 1992 (1992-01-08) the whole document, in particular claims 1 and 9	1-9
Y	WO 96 22300 A (MERCK & CO INC ; CVETOVICH RAYMOND (US); DEMCHAK RICHARD (US); MCCA) 25 July 1996 (1996-07-25) the whole document	1-9
	-/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

24 June 2002

Date of mailing of the international search report

03/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fitz, W

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/02044

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>MROZIK H ET AL:  "4-deoxy-4-aminoavermectins with potent  broad spectrum antiparasitic activities"  BIOORGANIC &amp; MEDICINAL CHEMISTRY LETTERS,  OXFORD, GB,  vol. 5, no. 20,  19 October 1995 (1995-10-19), pages  2435-2440, XP004135279  ISSN: 0960-894X  the whole document, in particular page  2438, table, entries 6-11  -----</p>	1-9
Y	<p>US 4 427 663 A (MROZIK HELMUT H)  24 January 1984 (1984-01-24)  cited in the application  the whole document, in particular column  20, line 35  -----</p>	1-9

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 02/02044

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5362863	A	08-11-1994	NONE	
EP 0465121	A	08-01-1992	AT 134377 T	15-03-1996
			AU 641777 B2	30-09-1993
			AU 7939291 A	02-01-1992
			CA 2045864 A1	29-12-1991
			DE 69117223 D1	28-03-1996
			DE 69117223 T2	12-09-1996
			DK 465121 T3	18-03-1996
			EP 0465121 A1	08-01-1992
			ES 2084110 T3	01-05-1996
			FI 913137 A ,B,	29-12-1991
			GR 3019067 T3	31-05-1996
			HK 157096 A	23-08-1996
			IE 912251 A1	01-01-1992
			IL 98599 A	29-06-1995
			JP 2112593 C	21-11-1996
			JP 5306294 A	19-11-1993
			JP 8022870 B	06-03-1996
			KR 224292 B1	15-10-1999
			LV 5740 A4	20-08-1996
			NO 178339 B	27-11-1995
			NO 952511 A	22-06-1995
			NZ 238705 A	27-07-1993
			PT 98117 A ,B	29-10-1993
			US 5288710 A	22-02-1994
			ZA 9104961 A	29-07-1992
WO 9622300	A	25-07-1996	AU 4699096 A	07-08-1996
			WO 9622300 A1	25-07-1996
US 4427663	A	24-01-1984	AU 556527 B2	06-11-1986
			AU 1239383 A	22-09-1983
			DE 3363686 D1	03-07-1986
			EP 0089202 A1	21-09-1983
			JP 1914207 C	23-03-1995
			JP 6033273 B	02-05-1994
			JP 58167591 A	03-10-1983
			MX 9202896 A1	30-06-1992
			NZ 203492 A	09-05-1986
			ZA 8301784 A	31-10-1984